



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/58	A1	(11) International Publication Number: WO 97/41873 (43) International Publication Date: 13 November 1997 (13.11.97)
(21) International Application Number: PCT/US97/07701 (22) International Filing Date: 6 May 1997 (06.05.97) (30) Priority Data: 60/017,377 7 May 1996 (07.05.96) US 60/019,336 6 June 1996 (06.06.96) US (71) Applicant: ERGO RESEARCH CORPORATION [US/US]; Unit #2, 212 Main Street, Wakefield, RI 02879 (US). (72) Inventor: CINCOTTA, Anthony, H.; 13 Shipway Place, Charlestown, MA 02129 (US). (74) Agents: LUDWIG, S., Peter et al.; Darby & Darby, P.C., 805 Third Avenue, New York, NY 10022 (US).	(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: METHOD AND COMPOSITION FOR THE TREATMENT OF LIPID AND GLUCOSE METABOLISM DISORDERS		
(57) Abstract <p>This invention is directed to modifying or regulating in a subject of at least one of lipid and glucose metabolism by administering to a subject in need of such treatment a dopamine D1 agonist in conjunction with one agent or agent combination selected from i) a dopamine D2 agonist, ii) at least one of an adrenergic alpha1 antagonist, an adrenergic alpha2 agonist and a serotonergic inhibitor, or iii) a dopamine D2 agonist further conjoined with at least one of an adrenergic alpha1 antagonist, an adrenergic alpha2 agonist and a serotonergic inhibitor.</p> <div data-bbox="1185 1848 1421 1921" style="text-align: right;">RECEIVED 11/15/97</div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

10

**METHOD AND COMPOSITION FOR THE TREATMENT OF
LIPID AND GLUCOSE METABOLISM DISORDERS**

This application claims priority under 35 U.S.C. § 119 from provisional
15 applications Serial Nos. 60/017,377 and 60/019,336, the disclosures of which are
incorporated herein in their entirety.

FIELD OF THE INVENTION

This invention relates to novel, improved methods for modifying or
20 regulating in a subject (vertebrate animal or human) of at least one of lipid and
glucose metabolism.

BACKGROUND OF THE INVENTION

Obesity and Lipid Metabolism Disorders - Body Fat Loss

In humans obesity can be defined as a body weight exceeding 20%
25 of the desirable body weight for individuals of the same sex, height and frame
(Salans, L.B., in Endocrinology & Metabolism, 2d Ed., McGraw-Hill, New York
1987, pp. 1203-1244; see also, R.H. Williams, Textbook of Endocrinology, 1974,
pp. 904-916). In other animals (or also in humans) obesity can be determined by
body weight patterns correlated with prolactin profiles given that members of a
30 species that are young, lean and "healthy" (i.e., free of any disorders, not just
metabolic disorders) have daily plasma prolactin level profiles that follow a pattern

characteristic of the species. This pattern is highly reproducible with a small standard deviation. Members of a species suffering from at least one of lipid and metabolism disorders, however, have aberrant prolactin profiles that depart from the normal (or healthy subjects') pattern by at least 1 SEM in at least two spaced
5 apart time points or by at least 2 SEM (standard error of the mean) in at least one time point.

Obesity, or excess fat deposits, correlate with and may trigger the onset of various lipid and/or glucose metabolism disorders, *e.g.* hypertension, Type II diabetes, atherosclerosis, *etc.*

10 Even in the absence of clinical obesity (according to the above definition) the reduction of body fat stores (notably visceral fat stores) in man especially on a long-term or permanent basis would be of significant benefit, both cosmetically and physiologically.

The reduction of body fat stores in domestic animals (as well as pets)
15 especially on a long-term or permanent basis would also obviously be of considerable economic benefit to man, particularly since farm animals supply a major portion of man's diet; and the animal fat may end up as *de novo* fat deposits in man.

Whereas controlled diet and exercise can produce modest results in
20 the reduction of body fat deposits, prior to the cumulative work of the present inventors (including the prior co-pending patent applications and issued U.S. patents referred to below), no truly effective or practical treatment had been found for controlling obesity or other lipid metabolism disorders.

Hyperlipoproteinemia is a condition in which the concentration of one
25 or more of cholesterol- or triglyceride-carrying lipoproteins (such as chylomicrons, very low density lipoproteins or VLDL and low-density lipoproteins or LDL) in plasma exceeds a normal limit. This upper limit is generally defined as the ninety-fifth percentile of a random population. Elevated levels of these substances have also been positively correlated with atherosclerosis and the often resulting cardiac
30 infarction, or "heart attack", which accounts for approximately half of all deaths in the United States. Strong clinical evidence has been presented which correlates a reduction in plasma lipoprotein concentration with a reduced risk of atherosclerosis (Noma, A., *et al.*, Atherosclerosis 49:1, 1983; Illingworth, D. and Conner, W.,

in Endocrinology & Metabolism, McGraw-Hill, New York 1987). Thus, a significant amount of research has been devoted to finding treatment methods which reduce levels of plasma cholesterol and triglycerides. High LDL and/or VLDL accompanied by high triglyceride levels in the blood constitute most important risk factors for atherosclerosis. Reduction of one or both of lipoproteins and triglycerides in the blood would reduce the risk of atherosclerosis and arrest or retard its development.

Another subset of the plasma lipoproteins found in vertebrates are high density lipoproteins, or HDL. HDL serve to remove free cholesterol from the plasma. A high HDL concentration as a percentage of total plasma cholesterol has been associated with a reduced risk of atherosclerosis and heart disease. Thus HDL are known in the lay press as "good" cholesterol. Therefore, therapeutic strategies involve attempts both to reduce plasma LDL and VLDL content (that is, reduce total plasma cholesterol), and to increase the HDL fraction of total plasma cholesterol. Several lines of research indicate that simply increasing HDL is of benefit even in the absence of LDL or VLDL reduction: Bell, G.P. *et al.*, Atherosclerosis 36:47-54, 1980; Fears, R., Biochem. Pharmacol. 33:219-228, 1984; Thompson, G., Br. Heart J. 51:585-588, 1989; Blackburn, H. N.E.J.M. 309:426-428, 1983.

Current therapies for hyperlipoproteinemias include a low fat diet and elimination of aggravating factors such as sedentary lifestyle. If the hyperlipoproteinemia is secondary (*i.e.* incident to *e.g.* a deficiency of lipoprotein lipase or LDL receptor, various endocrine pathologies, alcoholism, renal disorders, hepatic disorders) then control of the underlying disease is also central to treatment. Hyperlipoproteinemias are also treated with drugs, which usually alter the levels of particular components of the total plasma cholesterol, as well as reduce the total plasma lipid component. Among the most recently introduced drugs to treat hyperlipoproteinemia is lovastatin (MEVACOR®) which selectively inhibits an enzyme involved in cholesterol production, 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase. This drug specifically reduces total cholesterol and can cause a modest (5-10%) increase in HDL concentrations. However, benefits from these therapies vary from subject to subject.

Moreover, use of the HMG-CoA enzyme inhibitor is sometimes accompanied by side effects such as liver toxicity, renal myoglobinuria, renal

shutdown, and lenticular opacity. The risk of such side effects necessitates close monitoring of the patients (*e.g.*, liver function is tested monthly).

Another drug prescribed against hyperlipoproteinemia is clofibrate. The effectiveness of clofibrate also varies from subject to subject and its use is often accompanied by such side effects as nephrotic syndromes, myalgia, nausea and abdominal pain.

Diabetes and Glucose Metabolism Disorders

Diabetes, one of the most insidious of the major diseases, can strike suddenly or lie undiagnosed for years while attacking the blood vessels and nerves. Diabetics, as a group, are far more often afflicted with blindness, heart disease, stroke, kidney disease, hearing loss, gangrene and impotence. One third of all visits to physicians are occasioned by this disease and its complications, and diabetes and its complications are a leading cause of untimely death in the United States and in the Western world.

Diabetes adversely affects the way the body uses sugars and starches which, during digestion, are converted into glucose. Insulin, a hormone produced by the pancreas, makes the glucose available to the body's cells for energy. In muscle, adipose (fat) and connective tissues, insulin facilitates the entry of glucose into the cells by an action on the cell membranes. The ingested glucose is normally converted in the liver to CO₂ and H₂O (50%); to glycogen (5%); and to fat (30-40%), the latter being stored in fat depots. Fatty acids from the adipose tissues are circulated, returned to the liver for re-synthesis of triacylglycerol and metabolized to ketone bodies for utilization by the tissues. The fatty acids are also metabolized by other organs. Fat formation is a major pathway for carbohydrate utilization.

The net effect of insulin is to promote the storage and use of carbohydrates, protein and fat. Insulin deficiency is a common and serious pathologic condition in man. In insulin-dependent (IDDM or Type I) diabetes the pancreas produces little or no insulin, and insulin must be injected daily for the survival of the diabetic. In noninsulin-dependent (NIDDM or Type II) diabetes the pancreas retains the ability to produce insulin and in fact may produce higher than normal amounts of insulin, but the amount of insulin is relatively insufficient, or less than fully effective, due to cellular resistance to insulin.

In either form of diabetes there are widespread abnormalities. In most NIDDM subjects, the fundamental defects to which the abnormalities can be traced are (1) a reduced entry of glucose into various "peripheral" tissues and (2) an increased liberation of glucose into the circulation from the liver. There is therefore
5 an extracellular glucose excess and an intracellular glucose deficiency. There is also a decrease in the entry of amino acids into muscle and an increase in lipolysis. Hyperlipoproteinemia is also a complication of diabetes. The cumulative effect of these diabetes-associated abnormalities is severe blood vessel and nerve damage.

Other than the present invention and previous work by the present
10 inventors (discussed below), no effective treatment has been found for controlling either hyperinsulinemia or insulin resistance. Hyperinsulinemia is a higher-than-normal level of insulin in the blood. Insulin resistance can be defined as a state in which a normal amount of insulin produces a subnormal biologic response. In insulin-treated patients with diabetes, insulin resistance is considered to be present
15 whenever the therapeutic dose of insulin exceeds the secretory rate of insulin in normal persons. Insulin resistance is also associated with higher-than-normal levels of insulin *i.e.* hyperinsulinemia -- when normal or elevated levels of blood glucose are present.

20 Previous Work in This Field

Studies by the present inventors and others have indicated that the naturally occurring annual cycle of body fat store level, pervasive among vertebrates in the wild, reflects the activities of an adjustable central metabolostat that is comprised of circadian hypothalamic neural components. Changes in the
25 phase-relationships of circadian dopaminergic and serotonergic activities induce seasonal changes in metabolism and these circadian activities can be adjusted by appropriately timed treatments with hormones or neurotransmitter affecting drugs. In this regard, bromocriptine, a sympatholytic dopamine D₂ agonist with α_2 agonistic and α_1 antagonistic activities as well as serotonin inhibiting activities has
30 been demonstrated to reduce body fat store levels in a variety of animals including humans, without reducing food consumption, and also to reduce hyperinsulinemia, hyperlipidemia, and glucose intolerance.

The present inventors and their co-workers have previously found that administration of either or both of (i) certain prolactin reducing dopamine (D_2) agonists such as bromocriptine and (ii) prolactin-increasing substances such as dopamine antagonists, such as metoclopramide; and serotonin agonists and precursors, such as 5-hydroxytryptophan, reduce body fat stores, obesity, plasma triglycerides and cholesterol as well as hyperglycemia, hyperinsulinemia and insulin resistance: U.S. Patent Nos. 4,659,715; 4,749,709; 4,783,469; 5,006,526.

It is preferred to administer the prolactin reducing substances at a first predetermined time to effect a decrease in the circulating prolactin levels of the subject to be treated during an interval within the subject's daily prolactin cycle or rhythm when circulating (blood) prolactin levels are low in young, healthy subjects of the same species thereby causing the prolactin rhythm of the treated to approach or to conform to the standard or healthy subjects' prolactin rhythm. It is also preferred to administer the prolactin-increasing substances at a second predetermined time to effect an increase in the circulating prolactin levels of the subject to be treated during an interval within the subject's daily prolactin cycle or rhythm when circulating (blood) prolactin levels are high in young healthy subjects of the same species, thereby causing the prolactin rhythm of the treated subject to approach, or conform to, the standard or healthy subjects' prolactin rhythm. U.S. Patent Nos. 5,468,755; 5,496,803; 5,344,832, U.S. Patent 5,585,347 and U.S. Pat. Appln. 08/456,952 and PCT applications US93/12701 and US95/09061.

It is also known in the art that some of the effects of bromocriptine are supported by endogenous dopamine. (Ergot Compounds and Brain Functions Neuropsychiatric Aspects: Advances in Biochemical Psychopharmacology. M. Goldstein *et al.*, Eds. (Raven Press, New York, 1980) vol. 23). Specifically, it has been shown that locomotor stimulation and stereotyped behavior responses to bromocriptine are blocked by depletion of endogenous dopamine in rodents. However, if a D_1 agonist is subsequently provided to dopamine depleted animals, the responsiveness to bromocriptine is restored. Jackson, D.M. *et al.*, Psychopharmacology 94:321 (1988)). A similar dopaminergic $D_2:D_1$ interaction has been demonstrated in dopaminergic inhibition of feeding behavior. Although these studies confirm the importance of a $D_2:D_1$ interaction in the activation of dopaminergic activities, the increased locomotor activity and decreased feeding

response to $D_2:D_1$ agonists is acute and short lived, lasting for only a few hours. (Cooper, S.J. *et al.*, in $D_1:D_2$ Dopamine Receptor Interactions, J. Waddington, Ed. (Academic Press, London, 1993) pp. 203-234).

The previous work by third parties with D_1 and D_2 dopamine agonists in combination has not demonstrated any effects on lipid and glucose metabolism, and has not produced long-term responses of dopaminergic activities. Significantly, the present inventors have now found that the conjoined administration of a D_1 agonist and a D_2 dopamine agonist (or at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor) result in an unexpected and surprising improvement in one or more of the metabolic indices related to lipid and glucose metabolism when compared to the improvement (if any) provided by administration of a dopamine D_2 agonist such as bromocriptine administered alone.

15 OBJECTS OF THE INVENTION

It is one of the objects of this invention to provide additional improved methods for reducing in vertebrate subjects (including humans) in need of such treatment at least one of food consumption, body weight, body fat, plasma or blood glucose and blood insulin.

20 Another object of this invention is to provide methods for reducing at least one of insulin resistance (impaired glucose tolerance), hyperinsulinemia and hyperglycemia, and glycosylated hemoglobin (including A1C), and abating Type II diabetes.

A further object is to provide methods for reducing or retarding or 25 arresting atherosclerosis by reducing at least one of hyperlipoproteinemia and elevated blood triglycerides.

It is another object of this invention to provide methods for modifying and regulating lipid and glucose metabolism in a manner beneficial to the subject.

It is still another object of the invention to provide methods for 30 modifying and regulating lipid and glucose metabolism to provide effective treatments for obesity.

SUMMARY OF THE INVENTION

It has now been found that at least one of the foregoing objects can be accomplished by administering to a subject in need of such treatment a dopamine D₁ agonist in conjunction with one agent or agent combination selected
5 from the following:

- (i) a dopamine D₂ agonist;
- (ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor;
- (iii) a dopamine D₂ agonist further conjoined with at least one of an
10 adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor.

Preferably, the foregoing agents in (i), (ii) or (iii) above ("conjoined agents") are administered at a predetermined time i.e. within a restricted portion of a 24-hour period. Since the dopamine D₁ agonist amplifies the effect of the other agent or agents, the D₁ agonist is also preferably administered at about the
15 same time.

The conjoined administration of a dopamine D₁ agonist with one (or more) of the other agents identified above results in substantially augmented, and in fact often synergistic, effects in improvement of one or more metabolic indices related to glucose or lipid metabolism, and thus an improved modification or
20 regulation of at least one of glucose and lipid metabolism.

Where a D₂ agonist is employed, it is preferably an ergot alkaloid, most preferably bromocriptine.

In another aspect, the present invention is directed to administering to said subject:

- 25 (i) a D₂ agonist; and
- (ii) at least one agent, not a D₂ agonist, selected from the group consisting of adrenergic α_1 antagonists, adrenergic α_2 agonists and serotonergic inhibitors.

It has been found that such conjoined administration effects a greater
30 improvement on one or more of the foregoing metabolic indices than administering of a D₂ agonist singly.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 is a bar graph illustrating the weight loss (negative bars) or gain (positive bars) obtained in the experimental group administered both bromocriptine (BC) and SKF 38393 (SKF) compared to the animals administered SKF alone or BC alone or nothing (negative controls).

Fig. 2 is a graph of food intake (g/mouse/day) vs days of treatment of experimental ob/ob mice with both bromocriptine and SKF (dark circles) or no drug (open circles) or control lean animals given no drug (dark triangles).

Figs. 3A and 3B are bar graphs measuring fat body mass measured as glycerol (in g/mouse) (Fig. 3A) or lean body mass (protein in g/mouse) (Fig. 3B) for ob/ob animals that received no drug (control) or bromocriptine alone (second bar from left) or SKF alone (third bar from left) or both BC and SKF (fourth bar). The asterisk indicates a significant difference compared to the control bar.

Figs. 4A and 4B are bar graphs of blood glucose (mg/dl) of ob/ob animals (Fig. 4A) or serum insulin (ng/ml) of ob/ob animals (Fig. 4B) administered no drug (control); left most bar); BC alone (second bar from left); SKF alone (third bar from left) or both BC and SKF (fourth bar). The asterisks have the same significance as for Fig. 3A.

Figs. 5A and 5B are bar graphs of serum triglyceride levels (TG) in ng/dl (Fig. 5A) or serum free fatty acid levels (FFA) in mmol/l (Fig. 5B) for animals administered no drug (control; left most bar); BC alone (second bar from left); SKF alone (third bar from left) or both BC and SKF (fourth bar). The asterisks have the same significance as for Fig. 3A.

Figs. 6A-6C are bar graphs of blood glucose levels in mg/dl (Fig. 6A) serum triglyceride levels in mg/dl (Fig. 6B) and serum FFA in mmol/l (Fig. 6C) for animals administered no drug (left bar) or both BC and SKF (right bar). The asterisks have the same significance as for Fig. 3A. The animals were sacrificed at 3 HALO the lipogenesis peak for mice.

Figs. 7A-7C are bar graphs of liver enzyme activity (in millimoles of fatty acid per mg protein per minute) for the enzymes involved in fatty acid synthesis in the liver: fatty acid synthetase (Fig. 7A), malic enzyme (Fig. 7B) or glucose-6-phosphatase (Fig. 7C) illustrating difference in said activities as between

animals administered no drug (left bar) or both Bc and SKF (right bar). The asterisks have the same significance as for Fig. 3A.

Figs. 8A and 8B are bar graphs similar to those of Figs. 7A-7C but for the liver enzymes PEPCK (phosphoenol pyruvate carboxykinase) and glucose-6-phosphate dehydrogenase.

Figs. 9A-9C are bar graphs similar to those of Figs. 7A-7C but for the enzymes involved in fatty acid synthesis in adipose tissue: fatty acid synthetase (Fig. 9A) malic enzyme (Fig. 9B) and glucose-6-phosphate dehydrogenase (Fig. 9C).

Figs. 10A and 10B are bar graphs of glucose transport (in amoles/cell/minute) (Fig. 10A) and glucose oxidation in CO₂ (in amoles/cell/minute) (Fig. 10B) measured for BC + SKF treated and "no drug" mice in the absence (white bars) and presence (dark bars) of insulin in isolated adipocytes.

Fig. 11 is a bar graph of lipolysis measured as glycerol release (pmoles/cell/minute) in isolated adipocytes for BC + SKF treated and "no drug" mice.

Figs. 12A is a graph of adipose lipogenesis measured as rate of glycerol incorporation into lipids (mg/minute/gram of fat) as a function of the sacrifice time for mice (in HALO) treated with BC + SKF (open circles) or not treated (dark circles).

Fig. 12B is a bar graph of lipoprotein lipase (LPL) activity (in mmol of free fatty acid/10⁶ cells/hour for SKB + BC treated or "no drug" mice.

Figs. 13A and 13B are photomicrographs of adipocytes from BC + SKF treated (Fig. 13B) and untreated (Fig. 13A) animals. The amount of lipid per cell (in μ g lipid/cell) are given next to each Figure.

Figs. 14A-14C are photomicrographs of arcuate nuclei of ob/ob control mice (Fig. 14A) ob/ob BC + SKF treated mice (Fig. 14B) and lean (57 BL/6J) controls (Fig. 14C) showing large amounts of neuropeptide Y (NPY) mRNA in the ob/ob controls and significantly reduced amounts of NPY mRNA in the ob/ob treated mice.

Fig. 15 is a bar graph of NPY mRNA in the arcuate nucleus of ob/ob mice treated with BC + SKF (middle bar) or untreated ob/ob mice (left bar) or untreated lean controls (right bar).

Figure 16 is a plot of body weight v. day of treatment with a D₂ agonist alone or with D₁ agonist alone or with a combination of D₁/D₂ according to the invention. BC (10 mg/kg), BC plus SKF 38393, or vehicle injection on body weight in C57BL/6J ob/ob mice during two weeks of daily treatment at 1 hour after light onset. An asterisk denotes a significant difference in body weight change relative to all other treatment groups (P<0.02).

DETAILED DESCRIPTION OF THE INVENTION

All literature and patents and patent applications cited herein are incorporated by reference in their entirety. In case of a conflict, the present disclosure including its definitions shall control.

In one embodiment of the method of the present invention, a D₁ dopamine agonist is administered in conjunction with a second agent, consisting of at least one of a D₂ agonist, an α_2 agonist, an α_1 antagonist and a serotonergic inhibitor (or a D₂ agonist and at least one of the remaining agents) preferably at a specific time of day to a subject in need of treatment.

As used herein and applied to administration of more than one active ingredient the terms "conjoined" or "in conjunction" mean that the subject being thus treated receives a first active agent and also at least one other active agent, but not necessarily within the same formulation or dosage form and not necessarily at the same administration time. For example, the D₁ agonist and D₂ agonist or the other agent(s) can be administered at the same time (in the same dosage form or in two or more divided dosage forms) or sequentially at different times and in different dosage forms.

The D₁ dopamine agonist may be any one or more of those substances known to those skilled in the art that are capable of activating or potentiating D₁ dopamine receptors. The D₁ agonists that are suitable for use in the present invention include SKF38393, dihydrexidine, SKF 75670, SKF 82957, A77636, A68930, SKF 82526 (fenoldopam), and racemic trans-10, 11-dihydroxy 5, 6, 6a, 7, 8, 12b-hexahydro.

The D₂ agonists for use in the present invention can be any one or more of those compounds known to those skilled in the art that are capable of activating D₂ dopamine receptors. D₂ agonists suitable for use in the present

invention include LY-171555, bromocriptine methane sulfonate (+)-, 2,10,11-trihydroxyaporphine HBr, R(-)-, fisuride hydrogen maleate, 2-OH-NPA HCl, R(-)-, MDO-NPA HCl R(-), Propylnorpamorphine HCl R(-)-(NPA), and Quinperole HCl.

A preferred class of D_2 agonists includes ergot alkaloids such as
5 2-bromo-alpha-ergocriptine (bromocriptine), 6-methyl 8 beta -carbobenzyloxy-aminoethyl-10-alpha-ergoline, 8-acylaminoergoline, 6-methyl-8-alpha-(N-acyl)amino-9-ergoline, pergolide, lisuride, 6-methyl-8-alpha-(N-phenyl-acetyl)amino-9-ergoline, ergocornine, 9,10-dihydroergocornine, any D-2-halo-6-alkyl-8-substituted ergoline, and D-2-bromo-6-methyl-8-cyanomethylergoline. Of these bromocriptine is most
10 preferred.

Effective amounts of ergot alkaloid for humans and vertebrates when administered alone (not conjoined to a D_1 agonist) are typically within the range of 5.0 ug/kg/day to 0.2 mg/kg/day.

In general, effective amounts of D_2 agonist for humans and
15 vertebrates are within the range of 5ug/kg/day to 3.5 mg/kg/day.

The α_1 antagonists for use in the present invention can be any one or more of those compounds known to those skilled in the art that directly or indirectly block activation of α_1 adrenoceptors. The α_1 antagonists suitable for use in the present invention include bromocriptine, benoxathin HCl, naftopidil 2HCl,
20 (\pm)-niguldipine HCl, S(+)-niguldipine HCl, prazosin HCl, doxazosin HCl, spiperone HCl, urapidil HCl, 5-methyl urapidil, WB-4101 HCl.

Effective amounts of α_1 antagonist for humans and vertebrates are generally within the range of 0.02 to 0.3 mg/kg/day.

The α_2 agonists for use in the present invention can be any one or
25 more of those compounds known to those skilled in the art that are capable of activating α_2 adrenoceptors. The α_2 agonists suitable for use in the present invention include bromocriptine, agmatine sulfate, p-aminoclonidine HCl, B-HT 920 diHCl, B-HT 933 diHCl, clonidine HCl, guanabenz acetate, p-iodoclonidine HCl, oxymetazoline HCl, UK 14,304, and xylazine HCl.

30 Effective amounts of α_2 agonist for humans and vertebrates are generally within the range of 1 ug/kg/day to 0.3 mg/kg/day.

The serotonergic inhibitors suitable for use in the present invention include bromocriptine.

Effective amounts of serotonergic inhibitors for humans and vertebrates are generally within the range of 5 ug/kg/day to 0.2 mg/kg/day.

When two (or more) agents are administered in conjunction as disclosed in the Summary of Invention the amount of one or another can be lower
5 than stated above, and even amounts that are subthreshold (when an agent is used singly) can be employed.

The dopamine D₁ agonist and the dopamine D₂ agonist and/or other agent conjoined with the D₁ agonist (or with the D₂ agonist) may be administered to a subject preferably orally, or by subcutaneous, intravenous or intramuscular
10 injection. Dermal delivery systems, e.g., skin patches, as well as suppositories and other well-known systems for administering pharmaceutical agents can also be employed. Sublingual, nasal and other transmucosal modes of administration are also contemplated. Accelerated release compositions, such as those disclosed in U.S. Patent App'n Ser. No. 08/459,021, are preferred.

Each of the D₂ agonist, α_1 antagonist, α_2 agonist and serotonergic inhibitor are preferably administered at a predetermined time. The reason is that the effect of each of these agents on lipid and/or glucose metabolism is time-sensitive, as is explained in more detail for D₂ agonists in U.S. Patent 5,585,347 and U.S. Pat. Appln. 08/456,952, but applicable to the α_1 antagonists, α_2 agonists
20 and serotonergic inhibitors. The preferred time of administration is within an interval that results in effective blood levels of the agent(s) at a time during which the standard prolactin levels in healthy subjects of the species to be treated are low. For example in humans standard prolactin levels are low between the hours of 9:00 and 22:00. Accordingly, the predetermined time of administration of one
25 or more of the foregoing agents is between the hours of 5:00 and 13:00, preferably 7:00 and 12:00. Divided doses can be administered and the schedule of administration can be varied to take into account pharmacokinetic properties of each active agent. Details of administration are given in U.S. Patent 5,585,347 and U.S. Pat. Appln. 08/456,952 for bromocriptine, but also apply to the α_1
30 antagonists, α_2 agonists and serotonergic inhibitors employed in the present invention.

For mice the preferred time of administration of the active agent is within 1 hour after light onset. It is further preferred that the administration take place when the subject is neither active nor feeding.

For other vertebrate animals the preferred time of administration can be ascertained by reference to the standard prolactin rhythm for the species of the animal to be treated. The standard prolactin curve can be generated by measuring prolactin in young, healthy members of the species over a 24 hour period. See in U.S Patent 5,585,347 and U.S. Pat. Appln. 08/456,952.

The administration of the D_1 agonist is also preferably timed, i.e. the D_1 agonist is also administered at a predetermined time. Because the D_1 agonist amplifies the effect of the conjoined agent, it is advantageous to administer the D_1 agonist at or about the time of administration of the conjoined agent(s), such that the activity period of the D_1 agonist in the bloodstream of the treated subject overlaps (in fact preferably overlaps as much as possible) with the activity period of the conjoined agent. For convenience of administration and in order to promote subject compliance, the D_1 agonist can be administered at the same time as the conjoined agent(s).

The D_1 agonist may but need not be in the same formulation or dosage form (or form part of the same composition) as the conjoined agent(s). If more than one conjoined agent is administered, the conjoined agents may but need not be in the same formulation or dosage form or form part of the same composition.

In treating vertebrates, generally, dosages of the D_1 agonist and conjoined agent(s) are typically administered over a period ranging from about 10 days to about 180 days, or longer. Some patients (e.g., patients in particularly poor physical condition, or those of advanced age) may require a longer, or even continuous treatment. A treatment duration exceeding six months or even continuous treatment may be desirable even when not required.

At least one of body fat deposits, body weight, plasma or blood glucose, circulating insulin, plasma triglycerides (TG), plasma free fatty acids (FFA) and food consumption of the subject will be reduced as the result of the treatment. Disorders of lipid and glucose metabolism are thereby treated and subjects suffering from such pathologies as hyperphagia, obesity, insulin resistance

(impaired glucose tolerance), hyperlipidemia, hyperinsulinemia, and hyperglycemia will exhibit improvement in corresponding metabolic indices.

While appropriately timed administration of certain D_2 agonists (i.e., bromocriptine) alone will produce the effects described above to some degree, these effects are amplified (potentiated) by the conjoined administration of the D_1 agonist agents described in the present invention. In other words, the synergistic effect of the conjoined administration of the D_1 agonist and the conjoined agent (i.e., a D_2 agonist, and/or α_1 antagonist, and/or serotonergic inhibitor and/or α_2 agonist) produces results that are superior to those experienced through administration of the same amount of a D_2 agonist alone. It should be noted that the present invention permits but does not require each agent to be administered in an amount over the threshold amount (in the absence of a conjoined agent) to improve one or more metabolic indices precisely because of the augmented effect on these indices achieved by conjoined administration according to the present invention.

The benefits of the invention are not limited to modifying and regulating lipid and glucose metabolism. Other bodily functions, such as blood pressure, can be beneficially modified and regulated by timed administration of a D_2 agonist (as monotherapy) in the dosage range disclosed above. For example, the D_2 agonist bromocriptine administered at a dose within the range disclosed above (4.8 mg/day at 8:00 AM) has been shown by the present inventor to decrease significantly the diastolic blood pressure of humans. Conjoined administration of a dopamine D_1 agonist and

(i) a dopamine D_2 agonist;

(ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist, and a serotonergic inhibitor;

(iii) a dopamine D_2 agonist further conjoined with one or more of the members of (ii) above.

These and other features of the invention will be better understood by reference to the experiments described in the examples below.

EXAMPLE 1

Female ob/ob mice (40-70g bw) were treated for two weeks with either 1) bromocriptine (11 mg/kg) at light onset, 2) SKF38393 (20 mg/kg) at light onset, 3) bromocriptine plus SKF38393 at light onset, or 4) vehicle at light onset.

5 Bromocriptine or SKF38393 alone produced moderate reductions in hyperphagia, body weight gain, and obesity. However, bromocriptine plus SKF treatment produced significant reductions in hyperphagia (50-60% $p < 0.01$) resulting in dramatic weight loss (21%, $p < 0.0001$, compared to controls).

Body composition analysis of KOH/EtOH treated carcasses revealed
10 no significant decrease of protein mass and a 22% ($p < 0.05$) decrease of adipose mass in bromocriptine plus SKF treated mice relative to controls. Also, bromocriptine plus SKF treatment decreased to a much greater extent than bromocriptine or SKF alone, plasma free fatty acid (FFA) (44%, $p < 0.001$), triglyceride (TG) (50%, $p < 0.05$), and glucose (57%, $p < 0.01$). Insulin levels
15 tended to decrease (by 50%; $p < 0.09$) and total cholesterol remained unchanged by combined drug therapy.

Larger (65-75g) animals treated with bromocriptine plus SKF38393 demonstrated an even more dramatic loss of body weight relative to controls (10 + 1g in 10 days; $p < 0.01$). Arcuate neuropeptide Y (NPY) mRNA levels remain
20 unchanged after bromocriptine plus SKF treatment compared to controls.

C57BL/6J female obese mice of 40-45g body weight were treated by daily injections (at 1 HALO) of bromocriptine (BC at 10 mg/kg) and/or SKF38393 (SKF at 20 mg/kg). Animals were held on 12-hour daily photoperiods and fed ad libidum. Food consumption was monitored daily and body weights monitored at
25 days 0, 7 and 14 of the treatment.

Animals were sacrificed at 1 and/or 4 hours after light onset ("HALO") (except as described for Fig. 12A) and blood, liver and adipose tissue were collected. The carcasses were digested in ethanolic KOH and analyzed for protein and lipid content. Blood glucose was measured with an Accu-Chek Advantage
30 glucose meter (Boehringer). Serum insulin was measured with a radioimmunoassay kit (Linco Research) using rat insulin standards. Total triglycerides and free fatty acids were measured with kits from Sigma Diagnostics, St. Louis, MO and Wako Chemicals respectively.

Enzymatic activity of fatty acid synthase, malic enzyme and glucose-6-phosphate dehydrogenase was measured in isolated cytosolic fraction by spectrophotometric methods. Phosphoenolpyruvate carboxykinase (PEPCK) in liver cytosol was assayed by incorporation of $H^{14}CO_3^-$ into phosphoenolpyruvate.

- 5 Glucose-6-phosphatase activity was determined spectrophotometrically in isolated liver microsomes.

Adipocytes were isolated from perigonadal fat pads by collagenase digestion and their size was determined by combining microscopes measurement of their diameter and lipid extraction of their lipid content. Glucose transport and
10 glucose metabolism were measured by U- ^{14}C -glucose in the absence and presence of insulin and basal lipolysis was assayed by measuring glycerol release using a ^{32}P -g-ATP. Neuropeptide Y (NPY) mRNA levels were measured in the arcuate nuclei of the mice using in situ hybridization.

In summary, bromocriptine (BC) plus SKF38393 (SKF) treatment of
15 C57 BL/6 ob/ob mice produces the following changes in metabolic physiology:

- (1) A 42% reduction in hyperphagia, reducing daily feeding levels to less than or equal to lean (+/+) controls. (Fig. 2)
- (2) A 3.67g loss of body weight versus a 4.3g body weight gain in obese controls. (Fig. 1)
- 20 (3) A 27% reduction of body fat mass (Fig. 3A) with no loss of protein (Fig. 3B) despite substantial reduction in food consumption.
- (4) A 57 and 41% reduction in hyperglycemia (Fig. 4A) and hyperinsulinemia (Fig. 4B) respectively.
- (5) A 44 and 50% reduction serum FFA (Fig. 5B) and TG (Fig. 5A)
25 concentration.
- (6) A 27-78% reduction in lipogenesis enzymes within the liver and adipose (Figs. 7A-7C, 8A-8B, 9A-9C).
- (7) A 64% reduction in liver glucose-6-phosphatase and 80% increase in liver G6P dehydrogenase activities (Fig. 8B) as well as significant
30 reduction in fatty acid synthetase (Fig. 7A) and malic enzyme (Fig. 7B).
- (8) A 42% reduction in basal lipolysis from isolated adipocytes (Fig. 11) of in vivo treated mice with no change in glucose transport (Fig. 10A) or

oxidation (Fig. 10B) or GLUT4 expression (data not shown) as well as a significant reduction in adipocyte size (Compare Figs. 13B and 13A).

(9) A 50% reduction in adipose tissue lipoprotein lipase (LPL) activity and a blocking of lipogenesis (Figs. 12B and 12A respectively).

5 (10) The observed metabolic changes induced by BC + SKF are associated with a 30% reduction in NPY mRNA level within the arcuate nuclei resulting in levels still two fold greater than in lean (+ / +) counterparts (Fig. 15). A similar result can be qualitatively observed in Figs. 14A-14C.

(11) The reduction in blood glucose, triglyceride and free fatty acid
10 is more pronounced at 4 HALO (peak of lipolysis in the mouse). See Fig. 6 and compare Fig. 6A with Fig. 4A, Fig. 6B with Fig. 5A and Fig. 6C with Fig. 5B.

Treatment of genetically obese C57 BL/6J mice with bromocriptine (D₂ agonist) plus SKF38393 (D₁ agonist) induced a reduction of body weight associated with a marked (42%) reduction of hyperphagia. The resulting weight
15 loss was attributed nearly exclusively to loss of fat with protein mass remaining unchanged or even increased. Fat loss may be attributed to decreased caloric intake as well as decreased lipogenesis as both hepatic and adipose lipogenic enzyme activities were reduced by treatment. The substantial reduction in caloric intake induced by treatment was associated with a large reduction in circulating
20 free fatty acids (FFA). That is, fat cell size (lipid content) decreased appreciably while lipogenesis and lipid mobilization concurrently decreased. Apparently, the decreased mobilization is associated with an even greater decrease in lipid accretion. Such a conclusion is supported by the findings of decreased adipose LPL, and serum total and VLDL-TG (very low density lipoprotein/triglycerides).

25 The marked reduction in serum glucose induced by treatment is associated with a strong reduction in hepatic glucose-6-phosphatase activity and a somewhat less dramatic decrease in phosphoenolpyruvate carboxykinase activity. Interestingly, the reduction in hepatic G-6-phosphatase activity and the simultaneous increase in G-6-P dehydrogenase activity suggest specific metabolic
30 channelling towards glucose utilization in the liver rather than glucose release or production. Such alterations in liver metabolism facilitate increase of hepatic HADPH, nucleic acid, and protein synthesis.

The foregoing findings may be applied to treatment of humans suffering from obesity and other lipid disorders.

EXAMPLE 2

5 Different groups of 6-week old C57BL/6 ob/ob mice (lacking a functional leptin protein) were treated with either bromocriptine ("BC") (10mg/kg BW), SKF38393 ("SKF") (10mg/kg BW), both drugs, or vehicle for two weeks at 1 hour after light onset (HALO). Animals were held on 12-hour daily photoperiods and allowed to feed ad libitum. Food consumption was monitored daily for 3 days
10 before the initiation of treatment throughout the 14-day treatment period. Animals were sacrificed between 1 and 3 HALO on the day following the final treatment (i.e., 24-26 hours after last injection) and plasma was collected for the analyses of insulin, glucose, and lipids while the carcasses were solubilized in ethanolic KOH and analyzed for protein and lipid content. Bromocriptine and SKF38393,
15 individually, were ineffective in reducing body weight gain where as SKF, but not BC, reduced food consumption (19%, $P < 0.01$). However, the combined treatment of bromocriptine and SKF38393 (BC/SKF) decreased food consumption by 46% (from 4.8 ± 0.2 to 2.6 ± 0.2 g/day; $P < 0.001$) and body weight by 15% (from a 3.2g increase in controls to a 4.3g decrease; $P < 0.005$) in 14 days of treatment
20 (Fig. 16). Relative to controls, in absolute terms, the lipid content of the BC/SKF treated animals was decreased by 40% (from 4.2 ± 0.2 to 2.5 ± 0.3 g glycerol/animal; $P < 0.0003$) whereas the protein content increased 8% (from 3.7 ± 0.08 to 4.0 ± 0.08 g/animal; $P < 0.05$). Therefore, relative to control mice, the BC/SKF treated animals consumed less food but actually increased protein mass
25 while concurrently losing weight and fat. This effect on body composition was observed by SKF ($P < 0.003$) or BC ($P < 0.04$) treatments alone, although to a lesser extent than by BC/SKF combination ($P < 0.05$). Although BC alone and SKF alone significantly reduced plasma glucose concentration (by 31%; $P < 0.02$ and 43%; $P < 0.004$, respectively), the BC/SKF combination reduced plasma glucose (by
30 60%; $P < 0.0004$) substantially more than either drug alone ($P < 0.03$) to values equivalent to those values reported for lean euglycemic C57BL/6 mice (+/+)(1). Plasma insulin level was equally reduced by BC and BC/SKF treatment (50%; $P < 0.04$), but was not affected by SKF alone. BC/SKF, but neither BC nor SKF

individually, reduced plasma triglyceride and free fatty acid levels (by 36%; $P < 0.05$ and 44%; $P < 0.007$), (Table 1, below). These data indicate that the interactive effects of BC and SKF effectively reduced hyperphagia, obesity, insulin resistance, hyperglycemia, hyperinsulinemia, and hyperlipidemia in the ob/ob mouse.

Table 1. Effects of BC (10mg/kg), SKF (10mg/kg), BC plus SKF, or vehicle injections at 1 HALO on body weight, carcass composition, food consumption, and plasma glucose, insulin, and lipid levels of ob/ob mice following two weeks of treatment. Animals were sacrificed 24-26 hours following last treatment. Within parameters, values with similar superscripts denote a significant difference between treatments ($P < 0.05$ to < 0.0001).

	Final Body Weight (g)	Whole Body Lipid-Glycerol (g) (% BW)	Whole Body Protein (g) (% BW)	Plasma Glucose (mg/dl)	Plasma Insulin (ng/ml)	Plasma TG (mg/dl)	Plasma FFA (μ M)	Food Consumption (g/day)
CONTROL	64 ± 1^a	4.2 ± 0.2^a 7.9 ± 0.3^a	$3.7 \pm 0.1^{1,2,3}$ 6.9 ± 0.2^a	$380 \pm 39^{1,2}$	$69 \pm 12^{1,2}$	313 ± 49^a	826 ± 83^a	$48 \pm 0.2^{1,2}$
BC	63 ± 0.7^a	3.7 ± 0.1^a 7.0 ± 0.2^a	4.0 ± 0.1^a 7.5 ± 0.1^a	262 ± 25^a	30 ± 4^a	406 ± 101	818 ± 64^a	$4.6 \pm 0.2^{2,3}$
SKF	62 ± 0.7^a	3.1 ± 0.1^a 6.0 ± 0.2^a	4.1 ± 0.04^a 7.9 ± 0.1^a	218 ± 22^a	60 ± 13	234 ± 22^a	671 ± 36^a	$3.9 \pm 0.07^{1,2,3}$
BC/SKF	$46 \pm 2^{1,2,3}$	2.6 ± 0.3^a 5.4 ± 0.5^a	4.0 ± 0.1^a 6.9 ± 0.04^a	$164 \pm 15^{1,2}$	30 ± 9^a	199 ± 18	$461 \pm 83^{1,2,3}$	$2.6 \pm 0.2^{1,2,3}$

Values differ significantly from obese controls (a= $P < 0.05$; b= $P < 0.01$; c= $P < 0.001$).

EXAMPLE 3

The effects of BC/SKF treatment on circadian rhythms of key metabolic enzyme activities, serum metabolites and hormones regulating metabolism were examined. Obese C57BL/6J mice were treated for 2 weeks at 1 hour after light onset with BC (10 mg/kg BW) and SKF (20 mg/kg BW) or vehicle. Mice were then sacrificed every 4 hours over a 24 hr period for the analyses of serum hormones and metabolites and hepatic enzymatic activities. Serum glucose, free fatty acid (FFA) and hepatic glucose-6-phosphatase (G6Pase) activity were greatest during the light period of the day showing that this time period is the daily peak for lipolysis and hepatic glucose production in mice. BC/SKF treatment significantly reduced blood glucose (51%), FFA(56%) and G6Pase activity (38%) during this light period. Moreover, serum levels of the lipolytic and gluconeogenic hormones thyroxine and corticosterone were also highest during the light period and their levels were significantly reduced by 51%

and 53%, respectively by BC/SKF treatment. BC/SKF treatment also decreased the daily peak in liver phosphoenol pyruvate carboxykinase activity by 27% and increased the daily peak in liver glucose 6 phosphate dehydrogenase (by 32%) (potentiating glycolysis via xylose-5-phosphate production). Levels of serum insulin and liver malic enzyme were greatest during the dark period (feeding time) of the day illustrating increased lipogenesis during this time in mice. During this dark period BC/SKF treatment reduced serum insulin significantly, i.e., by 42%, and liver malic enzyme by 26%. BC/SKF treatment also decreased liver fatty acid synthase activity by 30-50%, normalizing its circadian rhythm. Such effects demonstrate the involvement both of circadian systems and BC/SKF influence on these systems in the regulation of metabolism.

EXAMPLE 4

The effect of *in vivo* BC/SKF treatment on glucose induced insulin release was studied *in vitro*. Obese (ob/ob) and lean (+ / +) C57BL/6J mice were treated daily for 2 weeks with BC (10 mg/kg) plus SKF (20 mg/kg) or vehicle only. Mice were sacrificed 25 hours after the final treatment and islets were isolated for static incubation with glucose. The BC/SKF treatment of obese mice reduced blood glucose (173 ± 14 mg/dl, $P < 0.01$), plasma total glycerol 162 ± 9 vs. 386 ± 33 mg/dl, $P < 0.01$), and plasma total cholesterol (143 ± 5 vs. 184 ± 5 mg/dl, $P < 0.01$) relative to obese controls. The plasma free fatty acid and insulin levels of treated mice were also reduced by 20-30% compared with that in obese controls. In control ob/ob mice, the insulin release from isolated islets stimulated by 10 mM glucose was the same as that by 8 mM glucose (1.6 ± 0.2 vs. 1.9 ± 0.5 ng/islet/h), while in BC/SKF treated ob/ob mice, 15 mM glucose induced a significant increase of insulin release compared with 8 mM glucose (4.1 ± 0.8 vs. 1.8 ± 0.4 ng/islet/h, $P < 0.05$). This enhancement is comparable to that observed in lean mice which exhibited a 2 fold increase of insulin release in response to 15 mM vs. 8 mM glucose. Similar BC/SKF treatment of lean mice showed no effect on glucose-stimulated insulin release from isolated islets compared to lean controls. BC/SKF treatment reversed impaired islet glucose sensing in ob/ob mice possibly due in part to the improvement of hyperglycemia and hyperlipidemia by this treatment.

Since hyperglycemia and hyperlipidemia may induce islet desensitization to glucose, which is a common syndrome in obesity-associated NIDDM in humans, the above finding can be applied to therapy of NIDDM in humans.

5

EXAMPLE 5

Metabolic changes resulting from the D₁/D₂ agonist treatment were evaluated in mice to determine if they were accompanied by decreases in density of NPY immunoreactivity in discrete hypothalamic nuclei. Female ob/ob mice (30-35g) were treated daily at 1 h after light onset with SKF38393 (20mg/kg) and bromocriptine (15mg/kg) or vehicle. Lean mice (C57BL/6J; 18-21g) treated with vehicle also served as controls. Following treatment for 12 days mice were sacrificed and their brains processed for NPY immunoreactivity. The treatment (summarized in Table 2 below) produced a significant decline in NPY levels in the SCN (38.5% P < 0.01), the arcuate nucleus (41%; P < 0.005) and the PVN (31.4% P < 0.05) compared to obese controls. In addition, during the study body weights increased in obese controls (8.3 ± 0.9g) whereas it decreased in treated animals (-1.1 ± 2g) (P < 0.0001). These results indicate that time of day-dependent dopaminergic D₁/D₂ coactivation improves hyperphagia, hyperglycemia and obesity in the ob/ob mouse, in part, by reducing elevated levels of hypothalamic NPY to that of lean animals.

Table 2

Type	Food consumed (g/day)	Blood glucose (mg/dl)	NPY density (arbitrary units)		
			SCN	Arcuate	PVN
Lean	3.1 ± 0.1	133 ± 5	39.8 ± 3	54 ± 4	49 ± 5
Obese	6.1 ± 0.1	216 ± 16	55.2 ± 4	95 ± 10	52 ± 6
Treated	4.3 ± 0.1 ^c	136 ± 9 ^c	34 ± 4 ^b	56 ± 8 ^b	36 ± 3 ^a

25

EXAMPLE 6

The influence of BC/SKF treatment on hepatic glucose metabolism was examined. Female C57BL/6J obese (ob/ob) mice (BW = 46 ± 1 g) were treated daily for 2 wks with BC (12.5 mg/kg) and SKF (20 mg/kg) or vehicle (n = 8-12/group) at 1 hr after light onset and then sacrificed at 24-26 hrs following the final day of treatment and liver tissue removed and analyzed for glucose 6-phosphatase (G6Pase) and glucose 6 phosphate dephydrogenase (G6PDase) activities and hepatic xylose-5-phosphate (X5P) concentration. Serum glucose and insulin levels were also determined. BC/SKF treatment significantly ($P < 0.01$) reduced serum glucose by 57% (from 435 ± 21 to 185 ± 8 mg/dl), serum insulin by 44% (from 25 ± 2 to 14 ± 3 ng/ml), hepatic G6Pase activity by 67% (from 1.5 ± 0.3 to 0.5 ± 0.07 μ moles/min/mg), and increased hepatic G6PDase activity by 73% (from 11 ± 1 to 19 ± 3 nmoles/min/mg), and X5P concentration by 73% (from 166 ± 10 to 287 ± 30 nmoles/g) relative to control. BC/SKF treatment resulted in a gluconeogenic substrate being shuttled away from glucose to the pentose phosphate pathway by the simultaneous inhibition of glucose 6-phosphatase (G6Pase) and stimulation of glucose- 6-phosphate dephydrogenase (G6PDase) thereby respectively blocking hepatic glucose production and shuttling glucose-6-phosphate towards production of xylose-5-phosphate (X5P), a potent activator of glycolysis. This is the first study to identify the existence of such a biochemical shift in hepatic glucose metabolism and its regulation by dopaminergic activation. Moreover, this dopaminergic regulated shift from hepatic gluconeogenesis towards potentiation of glycolysis could contribute to the normalization of severe hyperglycemia in these animals and may have significance in both the development and treatment of NIDDM in humans. Available evidence suggests that BC/SKF is acting in part at the ventromedial hypothalamus to produce these effects.

EXAMPLE 7

The combination of hyperglycemia and hypertriglyceridemia has been implicated as a risk factor for cardiovascular disease in NIDDM. Dopaminergic D_1/D_2 receptor co-activation with SKF38393 (SKF), a D_1 receptor agonist, plus bromocriptine (BC), a D_2 receptor agonist has been shown to act synergistically to reduce obesity. Its effects on hyperglycemia, dyslipidemia, and plasma lipoprotein

dynamics were tested in ob/ob mice. Obese C57BL/6J (ob/ob) mice (BW) 44.5 ± 0.5 g) were treated daily at light onset with vehicle (control) or SKF (20 mg/kg BW) plus BC (16 mg/kg BW) for 14 days. 25 to 28 hrs following the final treatment, animals were sacrificed and blood was collected for lipoprotein fractionation and analysis. Lipoprotein and serum triglyceride (TG), cholesterol (CH), phospholipid (PL), and serum glucose, insulin, and free fatty acid (FFA) were measured. White adipose, skeletal muscle, and heart tissues were harvested for analysis of lipoprotein lipase (LPL) activities. A second group of similarly treated animals was utilized for determination of hepatic triacylglycerol synthesis by following ^3H -glycerol incorporation into liver triglyceride 30 mins after its administration *in vivo*. 14 days of SKF/BC treatment significantly reduced blood glucose (390 ± 17 to 168 ± 6 mg/dl), serum TG (397 ± 22 to 153 ± 7 mg/dl), CH (178 ± 4 to 139 ± 4 mg/dl), PL (380 ± 7 to 263 ± 11 mg/dl), and FFA (1.1 ± 0.1 to 0.7 ± 0.1 mmol/l) ($P < 0.01$). Insulin was also reduced from 40 ± 5 to 28 ± 4 ng/ml ($P = 0.058$). Chylomicron-TG and VLDG-TG were reduced from 228 ± 2 to 45 ± 6 mg/dl and 169 ± 7 to 110 ± 4 mg/dl, respectively ($P < 0.01$). Hepatic triacylglycerol synthesis was reduced by 47% ($P < 0.01$). LPL activity was unchanged in skeletal and heart muscle tissues but was sharply reduced (67%) in adipose tissue ($P < 0.01$). LDL cholesterol level was reduced by 31% ($P < 0.01$). These data indicate that SKF/BC normalized hypertriglyceridemia via 1) decreasing Chylomicron-TG level and 2) decreasing VLDL-TG synthesis and secretion. The marked decrease in adipose LPL activity further supports the conclusion that serum VLDL-TG is reduced by decreased hepatic synthesis rather than increased removal from the circulation and may also contribute to the decreased serum FFA level. Moreover, the decreased serum FFA may contribute to the decreased hyperglycemia.

EXAMPLE 8

A 2-week treatment with SKF38393 (SKF), a dopamine D_1 receptor agonist, and bromocriptine (BC), a dopamine D_2 receptor agonist acts synergistically to reduce body fat and hyperglycemia in ob/ob mice in a food consumption independent manner. The biochemical mechanisms responsible for this effect were evaluated by measuring energy expenditure and metabolic

substrate utilization determined from respiratory quotient (RQ) of treated versus control mice. Circulating free fatty acid (FFA) levels represent the major rate limiting factor for fat oxidation and increased FFA also potentiate hyperglycemia in insulin resistant states. The influence of *in vivo* treatment with SKF38393 (SKF), a dopamine D₁ receptor agonist, and bromocriptine (BC), a dopamine D₂ receptor agonist, on serum FFA level and *in vitro* lipolysis in isolated adipocytes was tested. C57BL/6J obese (ob/ob) female mice were treated with vehicle (control) or SKF k(20 mg/kg BW) plus BC (10 mg/kg BW) for 14 days. BC/SKF treatment increased O₂ consumption and CO₂ production by 143% and 90% respectively (P<0.0001). Moreover, RQ values were shifted by treatment from 1.55±0.35 to 1.03±0.11 indicating a decrease in *de novo* glucose conversion to lipids (lipogenesis) and nearly exclusive utilization of glucose as an energy source (i.e., little fat oxidation). These findings are in accord with the substantial drug-induced decrease in serum glucose level (489±25 to 135±10 mg/dl, P<0.0001). These conclusions from the RQ data are further supported by a dramatic decrease in fat cell size (from 0.722±0.095 to 0.352±0.03 µg of lipid/cell, P<0.02) and a marked reduction in serum FFA levels (from 1.06±0.1 to 0.32±0.02 mM, P<0.001) and *in vitro* isoproterenol stimulated lipolysis (from 16.4±2.4 to 5±0.6 pmoles of glycerol released/cell/20 min, P<0.005). Therefore, the dramatic increase in O₂ and CO₂ production (and decreased fat cell size) cannot be explained by increased fat mobilization and oxidation. These data indicate that dopaminergic D₁,D₂ receptor coactivation shifts glucose metabolism from lipogenesis to oxidation with a concurrent decrease of fat mobilization and oxidation (thereby possibly improving insulin sensitivity). These findings have significance for the treatment of obesity and hypertriglyceridemia associated with NIDDM.

EXAMPLE 9

The combined effectiveness of SKF38393 (SKF), a D₁ receptor agonist, and bromocriptine (BC), a D₂ receptor agonist, were examined in treating obesity and diabetes in ob/ob (mice lacking the gene for the leptin protein) and db/db (mice lacking the gene for the leptin receptor) mice. Daily drug injections were administered to female C57BL/6J ob/ob and C57BL/KJ db/db mice 1 hr after light onset for 14 days. Drug treated groups received BC (16 mg/kg) plus SKF (20

mg/kg), whereas pair fed groups (food adjusted to drug treated groups' intake) and control groups received the vehicle. Oxygen consumption was measured in metabolic cages on day 11 or 12 of treatment. Plasma glucose, FFA, and insulin levels, were measured on day 14. In the ob/ob mice statistically significant results included: controls gained 6.9 ± 1.3 g of body weight, while the treated mice lost 7.4 ± 0.4 g. The average daily food consumption of controls was 6 ± 0.2 g versus 2.8 ± 0.1 g of treated. Oxygen consumption for controls and treated was 1277 ± 240 ml/kg/hr and 1623 ± 230 , respectively. Plasma glucose levels were 471 ± 42 mg/dl in controls, and 164 ± 13 in treated. k FFA levels were 1.27 ± 0.1 mM in controls, and 0.37 ± 0.05 in treated. Plasma insulin were 63.5 ± 17 ng/ml in controls, and 37.3 ± 6.6 in treated. Similar statistically significant results were observed in db/db mice: controls gained 6.6 ± 0.4 g, of body weight versus 3.4 ± 1.3 g in the treated. The average daily food consumption of controls was 10.7 ± 2.8 g versus 5.9 ± 0.5 g of the treated. Oxygen consumption for control and treated was 898 ± 2150 ml/kg/hr and 2322 ± 283 , respectively, Plasma glucose levels were 485 ± 29 mg/dl in controls, and 390 ± 55 in the treated. FFA levels were 1.49 ± 0.2 mM in controls, and 0.45 ± 0.04 in treated. Plasma from pairfed animals (in both ob/ob and db/db mice) indicate that the above drug-induced metabolic changes are not primarily the consequence of decreased food consumption. These results strongly suggest that hyperphagia, hyperglycemia and hyperlipidemia in animals lacking either leptin (ob/ob) or a functional leptin receptor (db/db) can be treated with the combined administration of D_1 and D_2 receptor agonists.

25 **EXAMPLE 10**

Pharmacological intervention with bromocriptine improves glucose and lipid metabolism in NIDDM animals and patients. The influence of such treatment on pancreatic islet function was investigated. The effect of D_1/D_2 receptor agonists -- bromocriptine/SKSF38393 (BC/SKF) on islet function in a mouse diabetic model was evaluated. Female db/db mice (30 ± 1 g) were treated daily for 2 weeks at 1 hr after light onset with 1) BC (16 mg/kg) plus SKF (20 mg/kg), 2) vehicle only (controls), or 3) vehicle plus feed restriction to match the reduced food consumption of treated mice (pair fed). The BC/SKF treatment reduced blood

glucose (347 ± 28 vs. 606 ± 31 mg/dl in controls, $P < 0.01$) and plasma free fatty acids (0.6 ± 0.1 vs. 1.1 ± 0.1 mM in controls, $P < 0.01$) levels, and increased plasma insulin level by 3-fold compared with that in controls (49 ± 5 vs. 16 ± 2 ng/ml, $P < 0.01$). In pair fed mice there was a more modest (30%) reduction (5 $P < 0.01$) of blood glucose but no change in plasma insulin and a 20% increase in plasma free fatty acids compared with control levels. The insulin release response of pancreatic islets to secretagogue was tested *in vitro*. Insulin release from incubated islets stimulated by glucose (8 and 15 mM), arginine (10 mM) and acetylcholine (10 μ M) was each 3-4 fold greater in the treated group compared 10 with that in controls ($P < 0.05$). Contrariwise, secretagogue-induced insulin release from incubated islets of pair fed mice were similar to those in controls. Furthermore, similar BC/SKF treatment had no effect in normal mice. Addition of BC/SKF directly to the islet incubation buffer did not enhance insulin release from db/db mouse islets. These results demonstrate the BC/SKF given *in vivo* markedly 15 enhance islet function in the db/db but not the normal mouse. This effect is not attributable to either a direct action on islet function or inhibition of feeding.

What is claimed is:

1. A method for modifying or regulating at least one of glucose or lipid metabolism disorders which comprises administering to a human or vertebrate animal subject in need of such modification or regulation a D₁ dopamine agonist in conjunction with a dopamine D₂ agonist wherein the conjoined administration is effective to improve at least one of the following lipid and glucose metabolic indices: body weight, body fat, plasma insulin, plasma glucose and plasma lipid, and plasma lipoprotein.

2. The method of claim 1 wherein the D₂ agonist is an ergot alkaloid is selected from the group consisting of 2-bromo-alpha-ergocryptive, 6-methyl 8 beta -carbobenzyloxyaminoethyl-10-alpha-ergoline, 8-acylaminoergoline, pergolide, lisuride, 6-methyl-8-alpha-(N-acyl)amino-9-ergoline, 6-methyl-8-alpha-(N-phenyl-acetyl)amino-9-ergoline, ergocomine, 9,10-dihydroergocomine, and D-2-halo-6-alkyl-8-substituted ergolines, D-2-bromo-6-methyl-8-cyanomethylergoline.

3. The method of claim 1 wherein the D₁ agonist comprises SKF38393.

4. The method of claim 2 wherein the ergot alkaloid comprises bromocriptine.

5. The method of claim 1 wherein the D₂ agonist is administered at a predetermined time of day.

6. The method of claim 5 wherein the D₁ agonist is administered at about the same time as the D₂ agonist.

7. A method for modifying or regulating at least one of glucose or lipid metabolism disorders which comprises administering to a human or vertebrate animal subject in need of such modification or regulation a D₁ agonist in

conjunction with at least one member selected from the group consisting of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor wherein the amount of the combination is sufficient to effect at least one of the following lipid and glucose metabolic indices: body weight, body fat, circulating insulin, plasma or blood glucose, plasma lipid and plasma lipoprotein.

8. The method of claim 7 wherein the D_1 agonist and the member of said group is administered at a predetermined time of day.

9. The method of claim 8 wherein the D_1 agonist is administered at about the same time as member of said group.

10. A method for modifying or regulating at least one of glucose or lipid metabolism disorders which comprises administering to a human or vertebrate animal subject in need of said modification or regulation a D_1 agonist in conjunction with at least one of a α_2 agonist, a α_1 antagonist and a serotonergic inhibitor, and further in conjunction with a D_2 agonist sufficient to improve at least one of the following lipid and glucose metabolic indices: body weight, body fat, plasma insulin, plasma glucose and plasma lipid.

11. A therapeutic agent combination for modifying or regulating at least one of glucose and lipid metabolism disorders comprising:

a first amount of a dopamine D_1 agonist; and
a second amount of at least one member selected from the group consisting of:

(i) a dopamine D_2 agonist; and
(ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor.

12. A method for modifying or regulating at least one of glucose or lipid metabolism disorders comprising:
administering to a subject in need of said modification or regulation:

- (i) a D_2 agonist in a first amount; and
 - (ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor in a second amount;
- said first and second amounts being sufficient to improve at least one of the
- 5 following indices of glucose or lipid metabolism: body weight, body fat, plasma insulin, plasma glucose and plasma lipid.

16 sheets of information
drawings as filed
4/28/97

Fig 1

Dramatic weight loss in BC+SKF group vs control
after 2 weeks of treatment






Fig. 2
BC+SKF treatment dramatically reduces
food intake in Ob/Ob mice





BC+SKF treatment selectively reduces fat body mass (by 27%) after 2 weeks of treatment

Fig. 3



BC+SKF treatment reduces blood glucose
and insulin levels by 40%

Fig. 4



Two weeks of BC+SKF treatment significantly reduces Triglyceride (by 50%) and Free Fatty Acid (by 44%) levels (Fig 5)




Fig 5A

Fig. 5B

6/16

At 4 hrs after the onset of lights (peak of lipolysis in the mouse) the reduction in blood glucose, triglyceride and free fatty acids is more pronounced




Fig 6A

Fig 6B

Fig 6C

BC+SKF treat. it reduces the activity of enzymes involved in fatty acid synthesis, gluconeogenesis and glucose production in the liver




Fig 7A




Fig 7B




Fig 7C

8/16

BC+SKF Reduction of enzyme activity in the liver

PEPCK




Fig. 8A

Glucose-6-Phosphate DH




Fig 8B

**BC+SKF treatment reduces the activity of
enzymes involved in fatty acid synthesis
in the adipose tissue**




Fig. 9A




Fig. 9B




Fig. 9C

BC+SKF treatment does not alter glucose transport or its oxydation in CO_2 in isolated adipocytes




Fig 10A





Fig 10B

Fig. 11

**BC+SKF treatment decreases lipolysis
in isolated adipocytes**



12/16

BC+SKF treatment blocks the nocturnal peak in adipose lipogenesis




Fig. 12A

BC+SKF treatment reduces LPL activity in the adipose tissue




Fig. 12B

Significant reduction in adipocyte size after
BC+SKF treatment Fig 13

control




Fig. 13A

μg lipid/cell

1.17 ± 0.05

BC+SKF




Fig. 13B

0.74 ± 0.03

Fig. 14
In situ hybridization for NPY
mRNA in the arcuate nuclei

Ob/Ob Control



Fig. 14A

Ob/Ob BC+SKF




Fig. 14B

Lean C57BL/6J Control




Fig. 14C

Fig. 15

NPY mRNA levels in the arcuate nucleus of obese mice decreased 31% after BC+SKF treatment but were still 2-fold higher compared to lean controls





Fig. 16



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/07701

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/58

US CL : 514/176, 172

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/176, 172

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database EMBASE on STN, EMBASE 90266123, GOLDSTEIN, M., 'Neurochemistry and psychopharmacology in self-injurious behavior: Dopaminergic mechanisms in self-inflicting biting behavior,' abstract, Psychopharmacol. Bull., 25/3, pp. 349-352, 1989, see entire abstract.	11

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 JULY 1997

Date of mailing of the international search report

05 SEP 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DWAYNE C. JONES

Telephone No. (703) 308-1235

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

EMBASE, HCAPLUS, MEDLINE, REGISTRY search terms: pergolide, bromocriptine, lisuride, ergocornine, ergoline, dopamine 1 receptor stimulating agent, dopamine 2 receptor stimulating agent, ergot alkal, alpha 1 adrenergic receptor blocking agent, alpha 2 adrenergic receptor stimulating agent, lipid metabolism or insulin metabolism, insulin blood level, glucose blood level, glucose brain level, lipid brain level, obesity, body weight, glucose, cholesterol, hypercholes, dopamine, lipid, insulin, metaboliz?, skf38393



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/58	A1	(11) International Publication Number: WO 97/41873 (43) International Publication Date: 13 November 1997 (13.11.97)
(21) International Application Number: PCT/US97/07701 (22) International Filing Date: 6 May 1997 (06.05.97) (30) Priority Data: 60/017,377 7 May 1996 (07.05.96) US 60/019,336 6 June 1996 (06.06.96) US (71) Applicant: ERGO RESEARCH CORPORATION [US/US]; Unit #2, 212 Main Street, Wakefield, RI 02879 (US). (72) Inventor: CINCOTTA, Anthony, H.; 13 Shipway Place, Charlestown, MA 02129 (US). (74) Agents: LUDWIG, S., Peter et al.; Darby & Darby, P.C., 805 Third Avenue, New York, NY 10022 (US).		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHOD AND COMPOSITION FOR THE TREATMENT OF LIPID AND GLUCOSE METABOLISM DISORDERS (57) Abstract <p>This invention is directed to modifying or regulating in a subject of at least one of lipid and glucose metabolism by administering to a subject in need of such treatment a dopamine D1 agonist in conjunction with one agent or agent combination selected from i) a dopamine D2 agonist, ii) at least one of an adrenergic alpha1 antagonist, an adrenergic alpha2 agonist and a serotonergic inhibitor, or iii) a dopamine D2 agonist further conjoined with at least one of an adrenergic alpha1 antagonist, an adrenergic alpha2 agonist and a serotonergic inhibitor.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BI	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

10

**METHOD AND COMPOSITION FOR THE TREATMENT OF
LIPID AND GLUCOSE METABOLISM DISORDERS**

This application claims priority under 35 U.S.C. § 119 from provisional
15 applications Serial Nos. 60/017,377 and 60/019,336, the disclosures of which are
incorporated herein in their entirety.

FIELD OF THE INVENTION

This invention relates to novel, improved methods for modifying or
20 regulating in a subject (vertebrate animal or human) of at least one of lipid and
glucose metabolism.

BACKGROUND OF THE INVENTION

Obesity and Lipid Metabolism Disorders - Body Fat Loss

In humans obesity can be defined as a body weight exceeding 20%
25 of the desirable body weight for individuals of the same sex, height and frame
(Salans, L.B., in Endocrinology & Metabolism, 2d Ed., McGraw-Hill, New York
1987, pp. 1203-1244; see also, R.H. Williams, Textbook of Endocrinology, 1974,
pp. 904-916). In other animals (or also in humans) obesity can be determined by
body weight patterns correlated with prolactin profiles given that members of a
30 species that are young, lean and "healthy" (i.e., free of any disorders, not just
metabolic disorders) have daily plasma prolactin level profiles that follow a pattern

characteristic of the species. This pattern is highly reproducible with a small standard deviation. Members of a species suffering from at least one of lipid and metabolism disorders, however, have aberrant prolactin profiles that depart from the normal (or healthy subjects') pattern by at least 1 SEM in at least two spaced
5 apart time points or by at least 2 SEM (standard error of the mean) in at least one time point.

Obesity, or excess fat deposits, correlate with and may trigger the onset of various lipid and/or glucose metabolism disorders, *e.g.* hypertension, Type II diabetes, atherosclerosis, *etc.*

10 Even in the absence of clinical obesity (according to the above definition) the reduction of body fat stores (notably visceral fat stores) in man especially on a long-term or permanent basis would be of significant benefit, both cosmetically and physiologically.

The reduction of body fat stores in domestic animals (as well as pets)
15 especially on a long-term or permanent basis would also obviously be of considerable economic benefit to man, particularly since farm animals supply a major portion of man's diet; and the animal fat may end up as *de novo* fat deposits in man.

Whereas controlled diet and exercise can produce modest results in
20 the reduction of body fat deposits, prior to the cumulative work of the present inventors (including the prior co-pending patent applications and issued U.S. patents referred to below), no truly effective or practical treatment had been found for controlling obesity or other lipid metabolism disorders.

Hyperlipoproteinemia is a condition in which the concentration of one
25 or more of cholesterol- or triglyceride-carrying lipoproteins (such as chylomicrons, very low density lipoproteins or VLDL and low-density lipoproteins or LDL) in plasma exceeds a normal limit. This upper limit is generally defined as the ninety-fifth percentile of a random population. Elevated levels of these substances have also been positively correlated with atherosclerosis and the often resulting cardiac
30 infarction, or "heart attack", which accounts for approximately half of all deaths in the United States. Strong clinical evidence has been presented which correlates a reduction in plasma lipoprotein concentration with a reduced risk of atherosclerosis (Noma, A., *et al.*, Atherosclerosis 49:1, 1983; Illingworth, D. and Conner, W.,

in Endocrinology & Metabolism, McGraw-Hill, New York 1987). Thus, a significant amount of research has been devoted to finding treatment methods which reduce levels of plasma cholesterol and triglycerides. High LDL and/or VLDL accompanied by high triglyceride levels in the blood constitute most important risk factors for atherosclerosis. Reduction of one or both of lipoproteins and triglycerides in the blood would reduce the risk of atherosclerosis and arrest or retard its development.

Another subset of the plasma lipoproteins found in vertebrates are high density lipoproteins, or HDL. HDL serve to remove free cholesterol from the plasma. A high HDL concentration as a percentage of total plasma cholesterol has been associated with a reduced risk of atherosclerosis and heart disease. Thus HDL are known in the lay press as "good" cholesterol. Therefore, therapeutic strategies involve attempts both to reduce plasma LDL and VLDL content (that is, reduce total plasma cholesterol), and to increase the HDL fraction of total plasma cholesterol. Several lines of research indicate that simply increasing HDL is of benefit even in the absence of LDL or VLDL reduction: Bell, G.P. *et al.*, Atherosclerosis 36:47-54, 1980; Fears, R., Biochem. Pharmacol. 33:219-228, 1984; Thompson, G., Br. Heart J. 51:585-588, 1989; Blackburn, H. N.E.J.M. 309:426-428, 1983.

Current therapies for hyperlipoproteinemias include a low fat diet and elimination of aggravating factors such as sedentary lifestyle. If the hyperlipoproteinemia is secondary (*i.e.* incident to *e.g.* a deficiency of lipoprotein lipase or LDL receptor, various endocrine pathologies, alcoholism, renal disorders, hepatic disorders) then control of the underlying disease is also central to treatment. Hyperlipoproteinemias are also treated with drugs, which usually alter the levels of particular components of the total plasma cholesterol, as well as reduce the total plasma lipid component. Among the most recently introduced drugs to treat hyperlipoproteinemia is lovastatin (MEVACOR®) which selectively inhibits an enzyme involved in cholesterol production, 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase. This drug specifically reduces total cholesterol and can cause a modest (5-10%) increase in HDL concentrations. However, benefits from these therapies vary from subject to subject.

Moreover, use of the HMG-CoA enzyme inhibitor is sometimes accompanied by side effects such as liver toxicity, renal myoglobinuria, renal

shutdown, and lenticular opacity. The risk of such side effects necessitates close monitoring of the patients (*e.g.*, liver function is tested monthly).

Another drug prescribed against hyperlipoproteinemia is clofibrate. The effectiveness of clofibrate also varies from subject to subject and its use is often accompanied by such side effects as nephrotic syndromes, myalgia, nausea and abdominal pain.

Diabetes and Glucose Metabolism Disorders

Diabetes, one of the most insidious of the major diseases, can strike suddenly or lie undiagnosed for years while attacking the blood vessels and nerves. Diabetics, as a group, are far more often afflicted with blindness, heart disease, stroke, kidney disease, hearing loss, gangrene and impotence. One third of all visits to physicians are occasioned by this disease and its complications, and diabetes and its complications are a leading cause of untimely death in the United States and in the Western world.

Diabetes adversely affects the way the body uses sugars and starches which, during digestion, are converted into glucose. Insulin, a hormone produced by the pancreas, makes the glucose available to the body's cells for energy. In muscle, adipose (fat) and connective tissues, insulin facilitates the entry of glucose into the cells by an action on the cell membranes. The ingested glucose is normally converted in the liver to CO₂ and H₂O (50%); to glycogen (5%); and to fat (30-40%), the latter being stored in fat depots. Fatty acids from the adipose tissues are circulated, returned to the liver for re-synthesis of triacylglycerol and metabolized to ketone bodies for utilization by the tissues. The fatty acids are also metabolized by other organs. Fat formation is a major pathway for carbohydrate utilization.

The net effect of insulin is to promote the storage and use of carbohydrates, protein and fat. Insulin deficiency is a common and serious pathologic condition in man. In insulin-dependent (IDDM or Type I) diabetes the pancreas produces little or no insulin, and insulin must be injected daily for the survival of the diabetic. In noninsulin-dependent (NIDDM or Type II) diabetes the pancreas retains the ability to produce insulin and in fact may produce higher than normal amounts of insulin, but the amount of insulin is relatively insufficient, or less than fully effective, due to cellular resistance to insulin.

In either form of diabetes there are widespread abnormalities. In most NIDDM subjects, the fundamental defects to which the abnormalities can be traced are (1) a reduced entry of glucose into various "peripheral" tissues and (2) an increased liberation of glucose into the circulation from the liver. There is therefore an extracellular glucose excess and an intracellular glucose deficiency. There is also a decrease in the entry of amino acids into muscle and an increase in lipolysis. Hyperlipoproteinemia is also a complication of diabetes. The cumulative effect of these diabetes-associated abnormalities is severe blood vessel and nerve damage.

Other than the present invention and previous work by the present inventors (discussed below), no effective treatment has been found for controlling either hyperinsulinemia or insulin resistance. Hyperinsulinemia is a higher-than-normal level of insulin in the blood. Insulin resistance can be defined as a state in which a normal amount of insulin produces a subnormal biologic response. In insulin-treated patients with diabetes, insulin resistance is considered to be present whenever the therapeutic dose of insulin exceeds the secretory rate of insulin in normal persons. Insulin resistance is also associated with higher-than-normal levels of insulin *i.e.* hyperinsulinemia -- when normal or elevated levels of blood glucose are present.

Previous Work in This Field

Studies by the present inventors and others have indicated that the naturally occurring annual cycle of body fat store level, pervasive among vertebrates in the wild, reflects the activities of an adjustable central metabolostat that is comprised of circadian hypothalamic neural components. Changes in the phase-relationships of circadian dopaminergic and serotonergic activities induce seasonal changes in metabolism and these circadian activities can be adjusted by appropriately timed treatments with hormones or neurotransmitter affecting drugs. In this regard, bromocriptine, a sympatholytic dopamine D_2 agonist with α_2 agonistic and α_1 antagonistic activities as well as serotonin inhibiting activities has been demonstrated to reduce body fat store levels in a variety of animals including humans, without reducing food consumption, and also to reduce hyperinsulinemia, hyperlipidemia, and glucose intolerance.

The present inventors and their co-workers have previously found that administration of either or both of (i) certain prolactin reducing dopamine agonists such as bromocriptine and (ii) prolactin-increasing substances such as dopamine antagonists, such as metoclopramide; and serotonin agonists and precursors, such as 5-hydroxytryptophan, reduce body fat stores, obesity, plasma triglycerides and cholesterol as well as hyperglycemia, hyperinsulinemia and insulin resistance: U.S. Patent Nos. 4,659,715; 4,749,709; 4,783,469; 5,006,526.

It is preferred to administer the prolactin reducing substances at a first predetermined time to effect a decrease in the circulating prolactin levels of the subject to be treated during an interval within the subject's daily prolactin cycle or rhythm when circulating (blood) prolactin levels are low in young, healthy subjects of the same species thereby causing the prolactin rhythm of the treated to approach or to conform to the standard or healthy subjects' prolactin rhythm. It is also preferred to administer the prolactin-increasing substances at a second predetermined time to effect an increase in the circulating prolactin levels of the subject to be treated during an interval within the subject's daily prolactin cycle or rhythm when circulating (blood) prolactin levels are high in young healthy subjects of the same species, thereby causing the prolactin rhythm of the treated subject to approach, or conform to, the standard or healthy subjects' prolactin rhythm. U.S. Patent Nos. 5,468,755; 5,496,803; 5,344,832, U.S. Patent 5,585,347 and U.S. Pat. Appln. 08/456,952 and PCT applications US93/12701 and US95/09061.

It is also known in the art that some of the effects of bromocriptine are supported by endogenous dopamine. (Ergot Compounds and Brain Functions Neuropsychiatric Aspects: Advances in Biochemical Psychopharmacology. M. Goldstein *et al.*, Eds. (Raven Press, New York, 1980) vol. 23). Specifically, it has been shown that locomotor stimulation and stereotyped behavior responses to bromocriptine are blocked by depletion of endogenous dopamine in rodents. However, if a D₁ agonist is subsequently provided to dopamine depleted animals, the responsiveness to bromocriptine is restored. Jackson, D.M. *et al.*, (Psychopharmacology 94:321 (1988)). A similar dopaminergic D₂:D₁ interaction has been demonstrated in dopaminergic inhibition of feeding behavior. Although these studies confirm the importance of a D₂:D₁ interaction in the activation of dopaminergic activities, the increased locomotor activity and decreased feeding

response to D₂:D₁ agonists is acute and short lived, lasting for only a few hours. (Cooper, S.J. *et al.*, in D₁:D₂ Dopamine Receptor Interactions, J. Waddington, Ed. (Academic Press, London, 1993) pp. 203-234).

The previous work by third parties with D₁ and D₂ dopamine agonists
5 in combination has not demonstrated any effects on lipid and glucose metabolism, and has not produced long-term responses of dopaminergic activities. Significantly, the present inventors have now found that the conjoined administration of a D₁ agonist and a D₂ dopamine agonist (or at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor)
10 result in an unexpected and surprising improvement in one or more of the metabolic indices related to lipid and glucose metabolism when compared to the improvement (if any) provided by administration of a dopamine D₂ agonist such as bromocriptine administered alone.

15 **OBJECTS OF THE INVENTION**

It is one of the objects of this invention to provide additional improved methods for reducing in vertebrate subjects (including humans) in need of such treatment at least one of food consumption, body weight, body fat, plasma or blood glucose and blood insulin.

20 Another object of this invention is to provide methods for reducing at least one of insulin resistance (impaired glucose tolerance), hyperinsulinemia and hyperglycemia, and glycosylated hemoglobin (including A1C), and abating Type II diabetes.

A further object is to provide methods for reducing or retarding or
25 arresting atherosclerosis by reducing at least one of hyperlipoproteinemia and elevated blood triglycerides.

It is another object of this invention to provide methods for modifying and regulating lipid and glucose metabolism in a manner beneficial to the subject.

It is still another object of the invention to provide methods for
30 modifying and regulating lipid and glucose metabolism to provide effective treatments for obesity.

SUMMARY OF THE INVENTION

It has now been found that at least one of the foregoing objects can be accomplished by administering to a subject in need of such treatment a dopamine D₁ agonist in conjunction with one agent or agent combination selected
5 from the following:

- (i) a dopamine D₂ agonist;
- (ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor;
- (iii) a dopamine D₂ agonist further conjoined with at least one of an
10 adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor.

Preferably, the foregoing agents in (i), (ii) or (iii) above ("conjoined agents") are administered at a predetermined time i.e. within a restricted portion of a 24-hour period. Since the dopamine D₁ agonist amplifies the effect of the other agent or agents, the D₁ agonist is also preferably administered at about the
15 same time.

The conjoined administration of a dopamine D₁ agonist with one (or more) of the other agents identified above results in substantially augmented, and in fact often synergistic, effects in improvement of one or more metabolic indices related to glucose or lipid metabolism, and thus an improved modification or
20 regulation of at least one of glucose and lipid metabolism.

Where a D₂ agonist is employed, it is preferably an ergot alkaloid, most preferably bromocriptine.

In another aspect, the present invention is directed to administering to said subject:

- 25 (i) a D₂ agonist; and
- (ii) at least one agent, not a D₂ agonist, selected from the group consisting of adrenergic α_1 antagonists, adrenergic α_2 agonists and serotonergic inhibitors.

It has been found that such conjoined administration effects a greater
30 improvement on one or more of the foregoing metabolic indices than administering of a D₂ agonist singly.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 is a bar graph illustrating the weight loss (negative bars) or gain (positive bars) obtained in the experimental group administered both bromocriptine (BC) and SKF 38393 (SKF) compared to the animals administered SKF alone or BC alone or nothing (negative controls).

Fig. 2 is a graph of food intake (g/mouse/day) vs days of treatment of experimental ob/ob mice with both bromocriptine and SKF (dark circles) or no drug (open circles) or control lean animals given no drug (dark triangles).

Figs. 3A and 3B are bar graphs measuring fat body mass measured as glycerol (in g/mouse) (Fig. 3A) or lean body mass (protein in g/mouse) (Fig. 3B) for ob/ob animals that received no drug (control) or bromocriptine alone (second bar from left) or SKF alone (third bar from left) or both BC and SKF (fourth bar). The asterisk indicates a significant difference compared to the control bar.

Figs. 4A and 4B are bar graphs of blood glucose (mg/dl) of ob/ob animals (Fig. 4A) or serum insulin (ng/ml) of ob/ob animals (Fig. 4B) administered no drug (control); left most bar); BC alone (second bar from left); SKF alone (third bar from left) or both BC and SKF (fourth bar). The asterisks have the same significance as for Fig. 3A.

Figs. 5A and 5B are bar graphs of serum triglyceride levels (TG) in ng/dl (Fig. 5A) or serum free fatty acid levels (FFA) in mmol/l (Fig. 5B) for animals administered no drug (control; left most bar); BC alone (second bar from left); SKF alone (third bar from left) or both BC and SKF (fourth bar). The asterisks have the same significance as for Fig. 3A.

Figs. 6A-6C are bar graphs of blood glucose levels in mg/dl (Fig. 6A) serum triglyceride levels in mg/dl (Fig. 6B) and serum FFA in mmol/l (Fig. 6C) for animals administered no drug (left bar) or both BC and SKF (right bar). The asterisks have the same significance as for Fig. 3A. The animals were sacrificed at 3 HALO the lipogenesis peak for mice.

Figs. 7A-7C are bar graphs of liver enzyme activity (in millimoles of fatty acid per mg protein per minute) for the enzymes involved in fatty acid synthesis in the liver: fatty acid synthetase (Fig. 7A), malic enzyme (Fig. 7B) or glucose-6-phosphatase (Fig. 7C) illustrating difference in said activities as between

animals administered no drug (left bar) or both Bc and SKF (right bar). The asterisks have the same significance as for Fig. 3A.

Figs. 8A and 8B are bar graphs similar to those of Figs. 7A-7C but for the liver enzymes PECK (phosphoenol pyruvate carboxykinase) and glucose-6-phosphate dehydrogenase.

Figs. 9A-9C are bar graphs similar to those of Figs. 7A-7C but for the enzymes involved in fatty acid synthesis in adipose tissue: fatty acid synthetase (Fig. 9A) malic enzyme (Fig. 9B) and glucose-6-phosphate dehydrogenase (Fig. 9C).

Figs. 10A and 10B are bar graphs of glucose transport (in amoles/cell/minute) (Fig. 10A) and glucose oxidation in CO_2 (in amoles/cell/minute) (Fig. 10B) measured for BC + SKF treated and "no drug" mice in the absence (white bars) and presence (dark bars) of insulin in isolated adipocytes.

Fig. 11 is a bar graph of lipolysis measured as glycerol release (pmoles/cell/minute) in isolated adipocytes for BC + SKF treated and "no drug" mice.

Figs. 12A is a graph of adipose lipogenesis measured as rate of glycerol incorporation into lipids (mg/minute/gram of fat) as a function of the sacrifice time for mice (in HALO) treated with BC + SKF (open circles) or not treated (dark circles).

Fig. 12B is a bar graph of lipoprotein lipase (LPL) activity (in mmol of free fatty acid/ 10^6 cells/hour for SKB + BC treated or "no drug" mice.

Figs. 13A and 13B are photomicrographs of adipocytes from BC + SKF treated (Fig. 13B) and untreated (Fig. 13A) animals. The amount of lipid per cell (in μg lipid/cell) are given next to each Figure.

Figs. 14A-14C are photomicrographs of arcuate nuclei of ob/ob control mice (Fig. 14A) ob/ob BC + SKF treated mice (Fig. 14B) and lean (57 BL/6J) controls (Fig. 14C) showing large amounts of neuropeptide Y (NPY) mRNA in the ob/ob controls and significantly reduced amounts of NPY mRNA in the ob/ob treated mice.

Fig. 15 is a bar graph of NPY mRNA in the arcuate nucleus of ob/ob mice treated with BC + SKF (middle bar) or untreated ob/ob mice (left bar) or untreated lean controls (right bar).

Figure 16 is a plot of body weight v. day of treatment with a D₂ agonist alone or with D₁ agonist alone or with a combination of D₁/D₂ according to the invention. BC (10 mg/kg), BC plus SKF 38393, or vehicle injection on body weight in C57BL/6J ob/ob mice during two weeks of daily treatment at 1 hour after light onset. An asterisk denotes a significant difference in body weight change relative to all other treatment groups (P < 0.02).

DETAILED DESCRIPTION OF THE INVENTION

All literature and patents and patent applications cited herein are incorporated by reference in their entirety. In case of a conflict, the present disclosure including its definitions shall control.

In one embodiment of the method of the present invention, a D₁ dopamine agonist is administered in conjunction with a second agent, consisting of at least one of a D₂ agonist, an α_2 agonist, an α_1 antagonist and a serotonergic inhibitor (or a D₂ agonist and at least one of the remaining agents) preferably at a specific time of day to a subject in need of treatment.

As used herein and applied to administration of more than one active ingredient the terms "conjoined" or "in conjunction" mean that the subject being thus treated receives a first active agent and also at least one other active agent, but not necessarily within the same formulation or dosage form and not necessarily at the same administration time. For example, the D₁ agonist and D₂ agonist or the other agent(s) can be administered at the same time (in the same dosage form or in two or more divided dosage forms) or sequentially at different times and in different dosage forms.

The D₁ dopamine agonist may be any one or more of those substances known to those skilled in the art that are capable of activating or potentiating D₁ dopamine receptors. The D₁ agonists that are suitable for use in the present invention include SKF38393, dihydrexidine, SKF 75670, SKF 82957, A77636, A68930, SKF 82526 (fenoldopam), and racemic trans-10, 11-dihydroxy 5, 6, 6a, 7, 8, 12b-hexahydro.

The D₂ agonists for use in the present invention can be any one or more of those compounds known to those skilled in the art that are capable of activating D₂ dopamine receptors. D₂ agonists suitable for use in the present

invention include LY-171555, bromocriptine methane sulfonate (+)-, 2,10,11-trihydroxyaporphine HBr, R(-)-, fisuride hydrogen maleate, 2-OH-NPA HCl, R(-)-, MDO-NPA HCl R(-), Propylorpamorphine HCl R(-)-(NPA), and Quinperole HCl.

A preferred class of D₂ agonists includes ergot alkaloids such as
5 2-bromo-alpha-ergocriptine (bromocriptine), 6-methyl 8 beta -carbobenzyloxy-aminoethyl-10-alpha-ergoline, 8-acylaminoergoline, 6-methyl-8-alpha-(N-acyl)amino-9-ergoline, pergolide, lisuride, 6-methyl-8-alpha-(N-phenyl-acetyl)amino-9-ergoline, ergocornine, 9,10-dihydroergocornine, any D-2-halo-6-alkyl-8-substituted ergoline, and D-2-bromo-6-methyl-8-cyanomethylergoline. Of these bromocriptine is most
10 preferred.

Effective amounts of ergot alkaloid for humans and vertebrates when administered alone (not conjoined to a D₁ agonist) are typically within the range of 5.0 ug/kg/day to 0.2 mg/kg/day.

In general, effective amounts of D₂ agonist for humans and
15 vertebrates are within the range of 5ug/kg/day to 3.5 mg/kg/day.

The α_1 antagonists for use in the present invention can be any one or more of those compounds known to those skilled in the art that directly or indirectly block activation of α_1 adrenoceptors. The α_1 antagonists suitable for use in the present invention include bromocriptine, benoxathin HCl, naftopidil 2HCl,
20 (\pm)-niguldipine HCl, S(+)-niguldipine HCl, prazosin HCl, doxazosin HCl, spiperone HCl, urapidil HCl, 5-methyl urapidil, WB-4101 HCl.

Effective amounts of α_1 antagonist for humans and vertebrates are generally within the range of 0.02 to 0.3 mg/kg/day.

The α_2 agonists for use in the present invention can be any one or
25 more of those compounds known to those skilled in the art that are capable of activating α_2 adrenoceptors. The α_2 agonists suitable for use in the present invention include bromocriptine, agmatine sulfate, p-aminoclonidine HCl, B-HT 920 diHCl, B-HT 933 diHCl, clonidine HCl, guanabenz acetate, p-iodoclonidine HCl, oxymetazoline HCl, UK 14,304, and xylazine HCl.

30 Effective amounts of α_2 agonist for humans and vertebrates are generally within the range of 1 ug/kg/day to 0.3 mg/kg/day.

The serotonergic inhibitors suitable for use in the present invention include bromocriptine.

Effective amounts of serotonergic inhibitors for humans and vertebrates are generally within the range of 5 ug/kg/day to 0.2 mg/kg/day.

When two (or more) agents are administered in conjunction as disclosed in the Summary of Invention the amount of one or another can be lower
5 than stated above, and even amounts that are subthreshold (when an agent is used singly) can be employed.

The dopamine D₁ agonist and the dopamine D₂ agonist and/or other agent conjoined with the D₁ agonist (or with the D₂ agonist) may be administered to a subject preferably orally, or by subcutaneous, intravenous or intramuscular
10 injection. Dermal delivery systems, e.g., skin patches, as well as suppositories and other well-known systems for administering pharmaceutical agents can also be employed. Sublingual, nasal and other transmucosal modes of administration are also contemplated. Accelerated release compositions, such as those disclosed in U.S. Patent App'n Ser. No. 08/459,021, are preferred.

Each of the D₂ agonist, α_1 antagonist, α_2 agonist and serotonergic
15 inhibitor are preferably administered at a predetermined time. The reason is that the effect of each of these agents on lipid and/or glucose metabolism is time-sensitive, as is explained in more detail for D₂ agonists in U.S. Patent 5,585,347 and U.S. Pat. Appln. 08/456,952, but applicable to the α_1 antagonists, α_2 agonists
20 and serotonergic inhibitors. The preferred time of administration is within an interval that results in effective blood levels of the agent(s) at a time during which the standard prolactin levels in healthy subjects of the species to be treated are low. For example in humans standard prolactin levels are low between the hours of 9:00 and 22:00. Accordingly, the predetermined time of administration of one
25 or more of the foregoing agents is between the hours of 5:00 and 13:00, preferably 7:00 and 12:00. Divided doses can be administered and the schedule of administration can be varied to take into account pharmacokinetic properties of each active agent. Details of administration are given in U.S. Patent 5,585,347 and U.S. Pat. Appln. 08/456,952 for bromocriptine, but also apply to the α_1
30 antagonists, α_2 agonists and serotonergic inhibitors employed in the present invention.

For mice the preferred time of administration of the active agent is within 1 hour after light onset. It is further preferred that the administration take place when the subject is neither active nor feeding.

For other vertebrate animals the preferred time of administration can be ascertained by reference to the standard prolactin rhythm for the species of the animal to be treated. The standard prolactin curve can be generated by measuring prolactin in young, healthy members of the species over a 24 hour period. See in U.S Patent 5,585,347 and U.S. Pat. Appln. 08/456,952.

The administration of the D_1 agonist is also preferably timed, i.e. the D_1 agonist is also administered at a predetermined time. Because the D_1 agonist amplifies the effect of the conjoined agent, it is advantageous to administer the D_1 agonist at or about the time of administration of the conjoined agent(s), such that the activity period of the D_1 agonist in the bloodstream of the treated subject overlaps (in fact preferably overlaps as much as possible) with the activity period of the conjoined agent. For convenience of administration and in order to promote subject compliance, the D_1 agonist can be administered at the same time as the conjoined agent(s).

The D_1 agonist may but need not be in the same formulation or dosage form (or form part of the same composition) as the conjoined agent(s). If more than one conjoined agent is administered, the conjoined agents may but need not be in the same formulation or dosage form or form part of the same composition.

In treating vertebrates, generally, dosages of the D_1 agonist and conjoined agent(s) are typically administered over a period ranging from about 10 days to about 180 days, or longer. Some patients (e.g., patients in particularly poor physical condition, or those of advanced age) may require a longer, or even continuous treatment. A treatment duration exceeding six months or even continuous treatment may be desirable even when not required.

At least one of body fat deposits, body weight, plasma or blood glucose, circulating insulin, plasma triglycerides (TG), plasma free fatty acids (FFA) and food consumption of the subject will be reduced as the result of the treatment. Disorders of lipid and glucose metabolism are thereby treated and subjects suffering from such pathologies as hyperphagia, obesity, insulin resistance

(impaired glucose tolerance), hyperlipidemia, hyperinsulinemia, and hyperglycemia will exhibit improvement in corresponding metabolic indices.

While appropriately timed administration of certain D₂ agonists (i.e., bromocriptine) alone will produce the effects described above to some degree, these effects are amplified (potentiated) by the conjoined administration of the D₁ agonist agents described in the present invention. In other words, the synergistic effect of the conjoined administration of the D₁ agonist and the conjoined agent (i.e., a D₂ agonist, and/or α_1 antagonist, and/or serotonergic inhibitor and/or α_2 agonist) produces results that are superior to those experienced through administration of the same amount of a D₂ agonist alone. It should be noted that the present invention permits but does not require each agent to be administered in an amount over the threshold amount (in the absence of a conjoined agent) to improve one or more metabolic indices precisely because of the augmented effect on these indices achieved by conjoined administration according to the present invention.

The benefits of the invention are not limited to modifying and regulating lipid and glucose metabolism. Other bodily functions, such as blood pressure, can be beneficially modified and regulated by timed administration of a D₂ agonist (as monotherapy) in the dosage range disclosed above. For example, the D₂ agonist bromocriptine administered at a dose within the range disclosed above (4.8 mg/day at 8:00 AM) has been shown by the present inventor to decrease significantly the diastolic blood pressure of humans. Conjoined administration of a dopamine D₁ agonist and

(i) a dopamine D₂ agonist;

(ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist, and a serotonergic inhibitor;

(iii) a dopamine D₂ agonist further conjoined with one or more of the members of (ii) above.

These and other features of the invention will be better understood by reference to the experiments described in the examples below.

EXAMPLE 1

Female ob/ob mice (40-70g bw) were treated for two weeks with either 1) bromocriptine (11 mg/kg) at light onset, 2) SKF38393 (20 mg/kg) at light onset, 3) bromocriptine plus SKF38393 at light onset, or 4) vehicle at light onset.

5 Bromocriptine or SKF38393 alone produced moderate reductions in hyperphagia, body weight gain, and obesity. However, bromocriptine plus SKF treatment produced significant reductions in hyperphagia (50-60% $p < 0.01$) resulting in dramatic weight loss (21%, $p < 0.0001$, compared to controls).

Body composition analysis of KOH/EtOH treated carcasses revealed
10 no significant decrease of protein mass and a 22% ($p < 0.05$) decrease of adipose mass in bromocriptine plus SKF treated mice relative to controls. Also, bromocriptine plus SKF treatment decreased to a much greater extent than bromocriptine or SKF alone, plasma free fatty acid (FFA) (44%, $p < 0.001$), triglyceride (TG) (50%, $p < 0.05$), and glucose (57%, $p < 0.01$). Insulin levels
15 tended to decrease (by 50%; $p < 0.09$) and total cholesterol remained unchanged by combined drug therapy.

Larger (65-75g) animals treated with bromocriptine plus SKF38393 demonstrated an even more dramatic loss of body weight relative to controls (10 + 1g in 10 days; $p < 0.01$). Arcuate neuropeptide Y (NPY) mRNA levels remain
20 unchanged after bromocriptine plus SKF treatment compared to controls.

C57BL/6J female obese mice of 40-45g body weight were treated by daily injections (at 1 HALO) of bromocriptine (BC at 10 mg/kg) and/or SKF38393 (SKF at 20 mg/kg). Animals were held on 12-hour daily photoperiods and fed ad libidum. Food consumption was monitored daily and body weights monitored at
25 days 0, 7 and 14 of the treatment.

Animals were sacrificed at 1 and/or 4 hours after light onset ("HALO") (except as described for Fig. 12A) and blood, liver and adipose tissue were collected. The carcasses were digested in ethanolic KOH and analyzed for protein and lipid content. Blood glucose was measured with an Accu-Chek Advantage
30 glucose meter (Boehringer). Serum insulin was measured with a radioimmunoassay kit (Linco Research) using rat insulin standards. Total triglycerides and free fatty acids were measured with kits from Sigma Diagnostics, St. Louis, MO and Wako Chemicals respectively.

Enzymatic activity of fatty acid synthase, malic enzyme and glucose-6-phosphate dehydrogenase was measured in isolated cytosolic fraction by spectrophotometric methods. Phosphoenolpyruvate carboxykinase (PEPCK) in liver cytosol was assayed by incorporation of $H^{14}CO_3^-$ into phosphoenolpyruvate.

- 5 Glucose-6-phosphatase activity was determined spectrophotometrically in isolated liver microsomes.

Adipocytes were isolated from perigonadal fat pads by collagenase digestion and their size was determined by combining microscopes measurement of their diameter and lipid extraction of their lipid content. Glucose transport and
10 glucose metabolism were measured by U- ^{14}C -glucose in the absence and presence of insulin and basal lipolysis was assayed by measuring glycerol release using a ^{32}P -g-ATP. Neuropeptide Y (NPY) mRNA levels were measured in the arcuate nuclei of the mice using in situ hybridization.

In summary, bromocriptine (BC) plus SKF38393 (SKF) treatment of
15 C57 BL/6 ob/ob mice produces the following changes in metabolic physiology:

- (1) A 42% reduction in hyperphagia, reducing daily feeding levels to less than or equal to lean (+/+) controls. (Fig. 2)
- (2) A 3.67g loss of body weight versus a 4.3g body weight gain in obese controls. (Fig. 1)
- 20 (3) A 27% reduction of body fat mass (Fig. 3A) with no loss of protein (Fig. 3B) despite substantial reduction in food consumption.
- (4) A 57 and 41% reduction in hyperglycemia (Fig. 4A) and hyperinsulinemia (Fig. 4B) respectively.
- (5) A 44 and 50% reduction serum FFA (Fig. 5B) and TG (Fig. 5A)
25 concentration.
- (6) A 27-78% reduction in lipogenesis enzymes within the liver and adipose (Figs. 7A-7C, 8A-8B, 9A-9C).
- (7) A 64% reduction in liver glucose-6-phosphatase and 80% increase in liver G6P dehydrogenase activities (Fig. 8B) as well as significant
30 reduction in fatty acid synthetase (Fig. 7A) and malic enzyme (Fig. 7B).
- (8) A 42% reduction in basal lipolysis from isolated adipocytes (Fig. 11) of in vivo treated mice with no change in glucose transport (Fig. 10A) or

oxidation (Fig. 10B) or GLUT4 expression (data not shown) as well as a significant reduction in adipocyte size (Compare Figs. 13B and 13A).

(9) A 50% reduction in adipose tissue lipoprotein lipase (LPL) activity and a blocking of lipogenesis (Figs. 12B and 12A respectively).

5 (10) The observed metabolic changes induced by BC + SKF are associated with a 30% reduction in NPY mRNA level within the arcuate nuclei resulting in levels still two fold greater than in lean (+/+) counterparts (Fig. 15). A similar result can be qualitatively observed in Figs. 14A-14C.

10 (11) The reduction in blood glucose, triglyceride and free fatty acid is more pronounced at 4 HALO (peak of lipolysis in the mouse). See Fig. 6 and compare Fig. 6A with Fig. 4A, Fig. 6B with Fig. 5A and Fig. 6C with Fig. 5B.

Treatment of genetically obese C57 BL/6J mice with bromocriptine (D₂ agonist) plus SKF38393 (D₁ agonist) induced a reduction of body weight associated with a marked (42%) reduction of hyperphagia. The resulting weight
15 loss was attributed nearly exclusively to loss of fat with protein mass remaining unchanged or even increased. Fat loss may be attributed to decreased caloric intake as well as decreased lipogenesis as both hepatic and adipose lipogenic enzyme activities were reduced by treatment. The substantial reduction in caloric intake induced by treatment was associated with a large reduction in circulating
20 free fatty acids (FFA). That is, fat cell size (lipid content) decreased appreciably while lipogenesis and lipid mobilization concurrently decreased. Apparently, the decreased mobilization is associated with an even greater decrease in lipid accretion. Such a conclusion is supported by the findings of decreased adipose LPL, and serum total and VLDL-TG (very low density lipoprotein/triglycerides).

25 The marked reduction in serum glucose induced by treatment is associated with a strong reduction in hepatic glucose-6-phosphatase activity and a somewhat less dramatic decrease in phosphoenolpyruvate carboxykinase activity. Interestingly, the reduction in hepatic G-6-phosphatase activity and the simultaneous increase in G-6-P dehydrogenase activity suggest specific metabolic
30 channelling towards glucose utilization in the liver rather than glucose release or production. Such alterations in liver metabolism facilitate increase of hepatic HADPH, nucleic acid, and protein synthesis.

The foregoing findings may be applied to treatment of humans suffering from obesity and other lipid disorders.

EXAMPLE 2

5 Different groups of 6-week old C57BL/6 ob/ob mice (lacking a functional leptin protein) were treated with either bromocriptine ("BC") (10mg/kg BW), SKF38393 ("SKF") (10mg/kg BW), both drugs, or vehicle for two weeks at 1 hour after light onset (HALO). Animals were held on 12-hour daily photoperiods and allowed to feed ad libitum. Food consumption was monitored daily for 3 days
10 before the initiation of treatment throughout the 14-day treatment period. Animals were sacrificed between 1 and 3 HALO on the day following the final treatment (i.e., 24-26 hours after last injection) and plasma was collected for the analyses of insulin, glucose, and lipids while the carcasses were solubilized in ethanolic KOH and analyzed for protein and lipid content. Bromocriptine and SKF38393,
15 individually, were ineffective in reducing body weight gain where as SKF, but not BC, reduced food consumption (19%, $P < 0.01$). However, the combined treatment of bromocriptine and SKF38393 (BC/SKF) decreased food consumption by 46% (from 4.8 ± 0.2 to 2.6 ± 0.2 g/day; $P < 0.001$) and body weight by 15% (from a 3.2g increase in controls to a 4.3g decrease; $P < 0.005$) in 14 days of treatment
20 (Fig. 16). Relative to controls, in absolute terms, the lipid content of the BC/SKF treated animals was decreased by 40% (from 4.2 ± 0.2 to 2.5 ± 0.3 g glycerol/animal; $P < 0.0003$) whereas the protein content increased 8% (from 3.7 ± 0.08 to 4.0 ± 0.08 g/animal; $P < 0.05$). Therefore, relative to control mice, the BC/SKF treated animals consumed less food but actually increased protein mass
25 while concurrently losing weight and fat. This effect on body composition was observed by SKF ($P < 0.003$) or BC ($P < 0.04$) treatments alone, although to a lesser extent than by BC/SKF combination ($P < 0.05$). Although BC alone and SKF alone significantly reduced plasma glucose concentration (by 31%; $P < 0.02$ and 43%; $P < 0.004$, respectively), the BC/SKF combination reduced plasma glucose (by
30 60%; $P < 0.0004$) substantially more than either drug alone ($P < 0.03$) to values equivalent to those values reported for lean euglycemic C57BL/6 mice (+/+)(1). Plasma insulin level was equally reduced by BC and BC/SKF treatment (50%; $P < 0.04$), but was not affected by SKF alone. BC/SKF, but neither BC nor SKF

individually, reduced plasma triglyceride and free fatty acid levels (by 36%; $P < 0.05$ and 44%; $P < 0.007$), (Table 1, below). These data indicate that the interactive effects of BC and SKF effectively reduced hyperphagia, obesity, insulin resistance, hyperglycemia, hyperinsulinemia, and hyperlipidemia in the ob/ob mouse.

Table 1. Effects of BC (10mg/kg), SKF (10mg/kg), BC plus SKF, or vehicle injections at 1 HALO on body weight, carcass composition, food consumption, and plasma glucose, insulin, and lipid levels of ob/ob mice following two weeks of treatment. Animals were sacrificed 24-26 hours following last treatment. Within parameters, values with similar superscripts denote a significant difference between treatments ($P < 0.05$ to < 0.0001).

	Final Body Weight (g)	Whole Body Lipid-Glycerol (g) (% BW)	Whole Body Protein (g) (% BW)	Plasma Glucose (mg/dl)	Plasma Insulin (ng/ml)	Plasma TG (mg/dl)	Plasma FFA (μ M)	Food Consumption (g/day)
CONTROL	54 \pm 1 ^a	4.2 \pm 0.2 ^a 7.8 \pm 0.3 ^a	3.7 \pm 0.1 ^{1,2,3} 6.9 \pm 0.2 ^a	380 \pm 39 ^{1,2}	59 \pm 12 ^{1,2}	313 \pm 49 ^a	826 \pm 83 ^a	4.9 \pm 0.2 ^{1,2}
BC	53 \pm 0.7 ^a	3.7 \pm 0.1 ^a 7.0 \pm 0.2 ^a	4.0 \pm 0.1 ^a 7.5 \pm 0.1 ^a	282 \pm 25 ^a	30 \pm 4 ^a	406 \pm 101 ^a	818 \pm 84 ^a	4.5 \pm 0.2 ^{2,3}
SKF	52 \pm 0.7 ^a	3.1 \pm 0.1 ^a 6.0 \pm 0.2 ^a	4.1 \pm 0.04 ^a 7.9 \pm 0.1 ^a	218 \pm 22 ^a	50 \pm 13 ^a	234 \pm 22 ^a	671 \pm 36 ^a	3.9 \pm 0.07 ^{3,4,5}
BC/SKF	45 \pm 2 ^{1,2,3}	2.5 \pm 0.3 ^a 5.4 \pm 0.5 ^a	4.0 \pm 0.1 ³ 8.8 \pm 0.04 ^a	154 \pm 15 ^{1,2}	30 \pm 9 ^a	199 \pm 18 ^a	461 \pm 63 ^{1,2,3}	2.6 \pm 0.2 ^{1,2,3}

Values differ significantly from obese controls (a = $P < 0.05$; b = $P < 0.01$; c = $P < 0.001$).

EXAMPLE 3

The effects of BC/SKF treatment on circadian rhythms of key metabolic enzyme activities, serum metabolites and hormones regulating metabolism were examined. Obese C57BL/6J mice were treated for 2 weeks at 1 hour after light onset with BC (10 mg/kg BW) and SKF (20 mg/kg BW) or vehicle. Mice were then sacrificed every 4 hours over a 24 hr period for the analyses of serum hormones and metabolites and hepatic enzymatic activities. Serum glucose, free fatty acid (FFA) and hepatic glucose-6-phosphatase (G6Pase) activity were greatest during the light period of the day showing that this time period is the daily peak for lipolysis and hepatic glucose production in mice. BC/SKF treatment significantly reduced blood glucose (51%), FFA(56%) and G6Pase activity (38%) during this light period. Moreover, serum levels of the lipolytic and gluconeogenic hormones thyroxine and corticosterone were also highest during the light period and their levels were significantly reduced by 51%

and 53%, respectively by BC/SKF treatment. BC/SKF treatment also decreased the daily peak in liver phosphoenol pyruvate carboxykinase activity by 27% and increased the daily peak in liver glucose 6 phosphate dehydrogenase (by 32%) (potentiating glycolysis via xylose-5-phosphate production). Levels of serum insulin and liver malic enzyme were greatest during the dark period (feeding time) of the day illustrating increased lipogenesis during this time in mice. During this dark period BC/SKF treatment reduced serum insulin significantly, i.e., by 42%, and liver malic enzyme by 26%. BC/SKF treatment also decreased liver fatty acid synthase activity by 30-50%, normalizing its circadian rhythm. Such effects demonstrate the involvement both of circadian systems and BC/SKF influence on these systems in the regulation of metabolism.

EXAMPLE 4

The effect of *in vivo* BC/SKF treatment on glucose induced insulin release was studied *in vitro*. Obese (ob/ob) and lean (+/+) C57BL/6J mice were treated daily for 2 weeks with BC (10 mg/kg) plus SKF (20 mg/kg) or vehicle only. Mice were sacrificed 25 hours after the final treatment and islets were isolated for static incubation with glucose. The BC/SKF treatment of obese mice reduced blood glucose (173 ± 14 mg/dl, $P < 0.01$), plasma total glycerol 162 ± 9 vs. 386 ± 33 mg/dl, $P < 0.01$), and plasma total cholesterol (143 ± 5 vs. 184 ± 5 mg/dl, $P < 0.01$) relative to obese controls. The plasma free fatty acid and insulin levels of treated mice were also reduced by 20-30% compared with that in obese controls. In control ob/ob mice, the insulin release from isolated islets stimulated by 10 mM glucose was the same as that by 8 mM glucose (1.6 ± 0.2 vs. 1.9 ± 0.5 ng/islet/h), while in BC/SKF treated ob/ob mice, 15 mM glucose induced a significant increase of insulin release compared with 8 mM glucose (4.1 ± 0.8 vs. 1.8 ± 0.4 ng/islet/h, $P < 0.05$). This enhancement is comparable to that observed in lean mice which exhibited a 2 fold increase of insulin release in response to 15 mM vs. 8 mM glucose. Similar BC/SKF treatment of lean mice showed no effect on glucose-stimulated insulin release from isolated islets compared to lean controls. BC/SKF treatment reversed impaired islet glucose sensing in ob/ob mice possibly due in part to the improvement of hyperglycemia and hyperlipidemia by this treatment.

Since hyperglycemia and hyperlipidemia may induce islet desensitization to glucose, which is a common syndrome in obesity-associated NIDDM in humans, the above finding can be applied to therapy of NIDDM in humans.

5

EXAMPLE 5

Metabolic changes resulting from the D₁/D₂ agonist treatment were evaluated in mice to determine if they were accompanied by decreases in density of NPY immunoreactivity in discrete hypothalamic nuclei. Female ob/ob mice (30-35g) were treated daily at 1 h after light onset with SKF38393 (20mg/kg) and bromocriptine (15mg/kg) or vehicle. Lean mice (C57BL/6J; 18-21g) treated with vehicle also served as controls. Following treatment for 12 days mice were sacrificed and their brains processed for NPY immunoreactivity. The treatment (summarized in Table 2 below) produced a significant decline in NPY levels in the SCN (38.5% P<0.01), the arcuate nucleus (41%; P<0.005) and the PVN (31.4% P<0.05) compared to obese controls. In addition, during the study body weights increased in obese controls (8.3 +/-0.9g) whereas it decreased in treated animals (-1.1 +/-2g) (P<0.0001). These results indicate that time of day-dependent dopaminergic D₁/D₂ coactivation improves hyperphagia, hyperglycemia and obesity in the ob/ob mouse, in part, by reducing elevated levels of hypothalamic NPY to that of lean animals.

Table 2

Type	Food consumed (g/day)	Blood glucose (mg/dl)	NPY density (arbitrary units)		
			SCN	Arcuate	PVN
Lean	3.1 +/-0.1	133 +/-5	39.8 +/-3	54 +/-4	49 +/-5
Obese	6.1 +/-0.1	216 +/-16	55.2 +/-4	95 +/-10	52 +/-6
Treated	4.3 +/-0.1 ^c	136 +/-9 ^c	34 +/-4 ^b	56 +/-8 ^b	36 +/-3 ^a

25

EXAMPLE 6

The influence of BC/SKF treatment on hepatic glucose metabolism was examined. Female C57BL/6J obese (ob/ob) mice (BW = 46 ± 1 g) were treated daily for 2 wks with BC (12.5 mg/kg) and SKF (20 mg/kg) or vehicle (n = 8-12/group) at 1 hr after light onset and then sacrificed at 24-26 hrs following the final day of treatment and liver tissue removed and analyzed for glucose 6-phosphatase (G6Pase) and glucose 6 phosphate dephydrogenase (G6PDase) activities and hepatic xylose-5-phosphate (X5P) concentration. Serum glucose and insulin levels were also determined. BC/SKF treatment significantly ($P < 0.01$) reduced serum glucose by 57% (from 435 ± 21 to 185 ± 8 mg/dl), serum insulin by 44% (from 25 ± 2 to 14 ± 3 ng/ml), hepatic G6Pase activity by 67% (from 1.5 ± 0.3 to 0.5 ± 0.07 μ moles/min/mg), and increased hepatic G6PDase activity by 73% (from 11 ± 1 to 19 ± 3 nmoles/min/mg), and X5P concentration by 73% (from 166 ± 10 to 287 ± 30 nmoles/g) relative to control. BC/SKF treatment resulted in a gluconeogenic substrate being shuttled away from glucose to the pentose phosphate pathway by the simultaneous inhibition of glucose 6-phosphatase (G6Pase) and stimulation of glucose- 6-phosphate dephydrogenase (G6PDase) thereby respectively blocking hepatic glucose production and shuttling glucose-6-phosphate towards production of xylose-5-phosphate (X5P), a potent activator of glycolysis. This is the first study to identify the existence of such a biochemical shift in hepatic glucose metabolism and its regulation by dopaminergic activation. Moreover, this dopaminergic regulated shift from hepatic gluconeogenesis towards potentiation of glycolysis could contribute to the normalization of severe hyperglycemia in these animals and may have significance in both the development and treatment of NIDDM in humans. Available evidence suggests that BC/SKF is acting in part at the ventromedial hypothalamus to produce these effects.

EXAMPLE 7

The combination of hyperglycemia and hypertriglyceridemia has been implicated as a risk factor for cardiovascular disease in NIDDM. Dopaminergic D_1/D_2 receptor co-activation with SKF38393 (SKF), a D_1 receptor agonist, plus bromocriptine (BC), a D_2 receptor agonist has been shown to act synergistically to reduce obesity. Its effects on hyperglycemia, dyslipidemia, and plasma lipoprotein

dynamics were tested in ob/ob mice. Obese C57BL/6J (ob/ob) mice (BW) 44.5 ± 0.5 g) were treated daily at light onset with vehicle (control) or SKF (20 mg/kg BW) plus BC (16 mg/kg BW) for 14 days. 25 to 28 hrs following the final treatment, animals were sacrificed and blood was collected for lipoprotein fractionation and analysis. Lipoprotein and serum triglyceride (TG), cholesterol (CH), phospholipid (PL), and serum glucose, insulin, and free fatty acid (FFA) were measured. White adipose, skeletal muscle, and heart tissues were harvested for analysis of lipoprotein lipase (LPL) activities. A second group of similarly treated animals was utilized for determination of hepatic triacylglycerol synthesis by following ^3H -glycerol incorporation into liver triglyceride 30 mins after its administration *in vivo*. 14 days of SKF/BC treatment significantly reduced blood glucose (390 ± 17 to 168 ± 6 mg/dl), serum TG (397 ± 22 to 153 ± 7 mg/dl), CH (178 ± 4 to 139 ± 4 mg/dl), PL (380 ± 7 to 263 ± 11 mg/dl), and FFA (1.1 ± 0.1 to 0.7 ± 0.1 mmol/l) ($P < 0.01$). Insulin was also reduced from 40 ± 5 to 28 ± 4 ng/ml ($P = 0.058$). Chylomicron-TG and VLDG-TG were reduced from 228 ± 2 to 45 ± 6 mg/dl and 169 ± 7 to 110 ± 4 mg/dl, respectively ($P < 0.01$). Hepatic triacylglycerol synthesis was reduced by 47% ($P < 0.01$). LPL activity was unchanged in skeletal and heart muscle tissues but was sharply reduced (67%) in adipose tissue ($P < 0.01$). LDL cholesterol level was reduced by 31% ($P < 0.01$). These data indicate that SKF/BC normalized hypertriglyceridemia via 1) decreasing Chylomicron-TG level and 2) decreasing VLDL-TG synthesis and secretion. The marked decrease in adipose LPL activity further supports the conclusion that serum VLDL-TG is reduced by decreased hepatic synthesis rather than increased removal from the circulation and may also contribute to the decreased serum FFA level. Moreover, the decreased serum FFA may contribute to the decreased hyperglycemia.

EXAMPLE 8

A 2-week treatment with SKF38393 (SKF), a dopamine D_1 receptor agonist, and bromocriptine (BC), a dopamine D_2 receptor agonist acts synergistically to reduce body fat and hyperglycemia in ob/ob mice in a food consumption independent manner. The biochemical mechanisms responsible for this effect were evaluated by measuring energy expenditure and metabolic

substrate utilization determined from respiratory quotient (RQ) of treated versus control mice. Circulating free fatty acid (FFA) levels represent the major rate limiting factor for fat oxidation and increased FFA also potentiate hyperglycemia in insulin resistant states. The influence of *in vivo* treatment with SKF38393 (SKF), a dopamine D₁ receptor agonist, and bromocriptine (BC), a dopamine D₂ receptor agonist, on serum FFA level and *in vitro* lipolysis in isolated adipocytes was tested. C57BL/6J obese (ob/ob) female mice were treated with vehicle (control) or SKF k(20 mg/kg BW) plus BC (10 mg/kg BW) for 14 days. BC/SKF treatment increased O₂ consumption and CO₂ production by 143% and 90% respectively (P<0.0001). Moreover, RQ values were shifted by treatment from 1.55±0.35 to 1.03±0.11 indicating a decrease in *de novo* glucose conversion to lipids (lipogenesis) and nearly exclusive utilization of glucose as an energy source (i.e., little fat oxidation). These findings are in accord with the substantial drug-induced decrease in serum glucose level (489±25 to 135±10 mg/dl, P<0.0001). These conclusions from the RQ data are further supported by a dramatic decrease in fat cell size (from 0.722±0.095 to 0.352±0.03 µg of lipid/cell, P<0.02) and a marked reduction in serum FFA levels (from 1.06±0.1 to 0.32±0.02 mM, P<0.001) and *in vitro* isoproterenol stimulated lipolysis (from 16.4±2.4 to 5±0.6 pmoles of glycerol released/cell/20 min, P<0.005). Therefore, the dramatic increase in O₂ and CO₂ production (and decreased fat cell size) cannot be explained by increased fat mobilization and oxidation. These data indicate that dopaminergic D₁/D₂ receptor coactivation shifts glucose metabolism from lipogenesis to oxidation with a concurrent decrease of fat mobilization and oxidation (thereby possibly improving insulin sensitivity). These findings have significance for the treatment of obesity and hypertriglyceridemia associated with NIDDM.

EXAMPLE 9

The combined effectiveness of SKF38393 (SKF), a D₁ receptor agonist, and bromocriptine (BC), a D₂ receptor agonist, were examined in treating obesity and diabetes in ob/ob (mice lacking the gene for the leptin protein) and db/db (mice lacking the gene for the leptin receptor) mice. Daily drug injections were administered to female C57BL/6J ob/ob and C57BL/KJ db/db mice 1 hr after light onset for 14 days. Drug treated groups received BC (16 mg/kg) plus SKF (20

mg/kg), whereas pair fed groups (food adjusted to drug treated groups' intake) and control groups received the vehicle. Oxygen consumption was measured in metabolic cages on day 11 or 12 of treatment. Plasma glucose, FFA, and insulin levels, were measured on day 14. In the ob/ob mice statistically significant results included: controls gained 6.9 ± 1.3 g of body weight, while the treated mice lost 7.4 ± 0.4 g. The average daily food consumption of controls was 6 ± 0.2 g versus 2.8 ± 0.1 g of treated. Oxygen consumption for controls and treated was 1277 ± 240 ml/kg/hr and 1623 ± 230 , respectively. Plasma glucose levels were 471 ± 42 mg/dl in controls, and 164 ± 13 in treated. k FFA levels were 1.27 ± 0.1 mM in controls, and 0.37 ± 0.05 in treated. Plasma insulin were 63.5 ± 17 ng/ml in controls, and 37.3 ± 6.6 in treated. Similar statistically significant results were observed in db/db mice: controls gained 6.6 ± 0.4 g, of body weight versus 3.4 ± 1.3 g in the treated. The average daily good consumption of controls was 10.7 ± 2.8 g versus 5.9 ± 0.5 g of the treated. Oxygen consumption for control and treated was 898 ± 2150 ml/kg/hr and 2322 ± 283 , respectively, Plasma glucose levels were 485 ± 29 mg/dl in controls, and 390 ± 55 in the treated. FFA levels were 1.49 ± 0.2 mM in controls, and 0.45 ± 0.04 in treated. Plasma from pairfed animals (in both ob/ob and db/db mice) indicate that the above drug-induced metabolic changes are not primarily the consequence of decreased food consumption. These results strongly suggest that hyperphagia, hyperglycemia and hyperlipidemia in animals lacking either leptin (ob/ob) or a functional leptin receptor (db/db) can be treated with the combined administration of D_1 and D_2 receptor agonists.

25 EXAMPLE 10

Pharmacological intervention with bromocriptine improves glucose and lipid metabolism in NIDDM animals and patients. The influence of such treatment on pancreatic islet function was investigated. The effect of D_1/D_2 receptor agonists -- bromocriptine/SKSF38393 (BC/SKF) on islet function in a mouse diabetic model was evaluated. Female db/db mice (30 ± 1 g) were treated daily for 2 weeks at 1 hr after light onset with 1) BC (16 mg/kg) plus SKF (20 mg/kg), 2) vehicle only (controls), or 3) vehicle plus feed restriction to match the reduced food consumption of treated mice (pair fed). The BC/SKF treatment reduced blood

glucose (347 ± 28 vs. 606 ± 31 mg/dl in controls, $P < 0.01$) and plasma free fatty acids (0.6 ± 0.1 vs. 1.1 ± 0.1 mM in controls, $P < 0.01$) levels, and increased plasma insulin level by 3-fold compared with that in controls (49 ± 5 vs. 16 ± 2 ng/ml, $P < 0.01$). In pair fed mice there was a more modest (30%) reduction (5 $P < 0.01$) of blood glucose but no change in plasma insulin and a 20% increase in plasma free fatty acids compared with control levels. The insulin release response of pancreatic islets to secretagogue was tested *in vitro*. Insulin release from incubated islets stimulated by glucose (8 and 15 mM), arginine (10 mM) and acetylcholine (10 μ M) was each 3-4 fold greater in the treated group compared 10 with that in controls ($P < 0.05$). Contrariwise, secretagogue-induced insulin release from incubated islets of pair fed mice were similar to those in controls. Furthermore, similar BC/SKF treatment had no effect in normal mice. Addition of BC/SKF directly to the islet incubation buffer did not enhance insulin release from db/db mouse islets. These results demonstrate the BC/SKF given *in vivo* markedly 15 enhance islet function in the db/db but not the normal mouse. This effect is not attributable to either a direct action on islet function or inhibition of feeding.

What is claimed is:

1. A method for modifying or regulating at least one of glucose or lipid metabolism disorders which comprises administering to a human or vertebrate animal subject in need of such modification or regulation a D₁ dopamine agonist in conjunction with a dopamine D₂ agonist wherein the conjoined administration is effective to improve at least one of the following lipid and glucose metabolic indices: body weight, body fat, plasma insulin, plasma glucose and plasma lipid, and plasma lipoprotein.
2. The method of claim 1 wherein the D₂ agonist is an ergot alkaloid is selected from the group consisting of 2-bromo-alpha-ergocriptine, 6-methyl 8 beta -carbobenzyloxyaminoethyl-10-alpha-ergoline, 8-acylaminoergoline, pergolide, lisuride, 6-methyl-8-alpha-(N-acyl)amino-9-ergoline, 6-methyl-8-alpha-(N-phenyl-acetyl)amino-9-ergoline, ergocomine, 9,10-dihydroergocomine, and D-2-halo-6-alkyl-8-substituted ergolines, D-2-bromo-6-methyl-8-cyanomethylergoline.
3. The method of claim 1 wherein the D₁ agonist comprises SKF38393.
4. The method of claim 2 wherein the ergot alkaloid comprises bromocriptine.
5. The method of claim 1 wherein the D₂ agonist is administered at a predetermined time of day.
6. The method of claim 5 wherein the D₁ agonist is administered at about the same time as the D₂ agonist.
7. A method for modifying or regulating at least one of glucose or lipid metabolism disorders which comprises administering to a human or vertebrate animal subject in need of such modification or regulation a D₁ agonist in

conjunction with at least one member selected from the group consisting of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor wherein the amount of the combination is sufficient to effect at least one of the following lipid and glucose metabolic indices: body weight, body fat, circulating insulin, plasma or blood glucose, plasma lipid and plasma lipoprotein.

8. The method of claim 7 wherein the D_1 agonist and the member of said group is administered at a predetermined time of day.

9. The method of claim 8 wherein the D_1 agonist is administered at about the same time as member of said group.

10. A method for modifying or regulating at least one of glucose or lipid metabolism disorders which comprises administering to a human or vertebrate animal subject in need of said modification or regulation a D_1 agonist in conjunction with at least one of a α_2 agonist, a α_1 antagonist and a serotonergic inhibitor, and further in conjunction with a D_2 agonist sufficient to improve at least one of the following lipid and glucose metabolic indices: body weight, body fat, plasma insulin, plasma glucose and plasma lipid.

11. A therapeutic agent combination for modifying or regulating at least one of glucose and lipid metabolism disorders comprising:

a first amount of a dopamine D_1 agonist; and
a second amount of at least one member selected from the group consisting of:

- (i) a dopamine D_2 agonist; and
- (ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor.

12. A method for modifying or regulating at least one of glucose or lipid metabolism disorders comprising:
administering to a subject in need of said modification or regulation:

- (i) a D_2 agonist in a first amount; and
 - (ii) at least one of an adrenergic α_1 antagonist, an adrenergic α_2 agonist and a serotonergic inhibitor in a second amount;
- said first and second amounts being sufficient to improve at least one of the
- 5 following indices of glucose or lipid metabolism: body weight, body fat, plasma insulin, plasma glucose and plasma lipid.

*16 sheets of informal
drawings as filed
4/28/97*

Fig 1

**Dramatic weight loss in BC+SKF group vs control
after 2 weeks of treatment**






Fig. 2
BC+SKF treatment dramatically reduces
food intake in Ob/Ob mice




BC+SKF treatment selective reduces fat body mass (by 27%) after 2 weeks of treatment

Fig. 3

Fat Body Mass





Lean Body Mass



BC+SKF treatment reduces blood glucose
and insulin levels by 40%

Fig. 4



Two weeks of BC+SKF treatment significantly reduces Triglyceride (by 50%) and Free Fatty Acid (by 44%) levels

(Fig 5)




Fig 5A

Fig. 5B

6/16

At 4 hrs after the onset of lights (peak of lipolysis in the mouse) the reduction in blood glucose, triglyceride and free fatty acids is more pronounced




Fig 6A

Fig 6B

Fig 6C

BC+SKF treat. it reduces the activity of enzymes involved in fatty acid synthesis, gluconeogenesis and glucose production in the liver




Fig 7A




Fig 7B




Fig 7C

8/16

BC+SKF Reduction of enzyme actⁿ in the liver
PEPCK




Fig. 8A

Glucose-6-Phosphate DH




Fig 8B

**BC+SKF treatment reduces the activity of
enzymes involved in fatty acid synthesis
in the adipose tissue**




Fig. 9A




Fig. 9B




Fig. 9C

BC+SKF treatment does not alter glucose transport or its oxydation in CO_2 in isolated adipocytes




Fig 10A





Fig 10B

Fig. 11

**BC+SKF treatment decreases lipolysis
in isolated adipocytes**

12/16

BC+SKF treatment blocks the nocturnal peak in adipose lipogenesis




Fig. 12A

BC+SKF treatment reduces LPL activity in the adipose tissue




Fig. 12B

Significant reduction in adipocyte size after
BC+SKF treatment Fig 13

control




Fig. 13A

μg lipid/cell

1.17 ± 0.05

BC+SKF




Fig. 13B

0.74 ± 0.03

Fig. 14
In situ hybridization for NPY
mRNA in the arcuate nuclei

Ob/Ob Control



Fig. 14A

Ob/Ob BC+SKF



Fig. 14B

Lean C57BL/6J Control



Fig. 14C

Fig. 15

NPY mRNA levels in the arcuate nucleus of obese mice decreased 31% after BC+SKF treatment but were still 2-fold higher compared to lean controls





FIG. 16



A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/58

US CL : 514/176, 172

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/176, 172

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

None

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database EMBASE on STN, EMBASE 90266123, GOLDSTEIN, M., 'Neurochemistry and psychopharmacology in self-injurious behavior: Dopaminergic mechanisms in self-inflicting biting behavior,' abstract, Psychopharmacol. Bull., 25/3, pp. 349-352, 1989, see entire abstract.	11

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	documents of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

21 JULY 1997

Date of mailing of the international search report

05 SEP 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DWAYNE C. JONES

Telephone No. (703) 308-1235

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

EMBASE, HCAPLUS, MEDLINE, REGISTRY search terms: pergolide, bromocriptine, lisuride, egrocoornine, ergoline, dopamine 1 receptor stimulating agent, dopamine 2 receptor stimulating agent, ergot alkal, alpha 1 adrenergic receptor blocking agent, alpha 2 adrenergic receptor stimulating agent, lipid metabolism or insulin metabolism, insulin blood level, glucose blood level, glucose brain level, lipid brain level, obesity, body weight, glucose, cholesterol, hypercholes, dopamine, lipid, insulin, metaboli?, skf38393